

Chitosans as bioactive macromolecules to protect economically relevant crops from their main pathogens

✉ Alejandro Falcón¹, Aida T Rodríguez², Miguel A Ramírez², Deyanira Rivero², Benedicto Martínez³, Juan C Cabrera⁴, Daimy Costales¹, Ariel Cruz², Luis G González⁵, María C Jiménez⁵, Leonel Jiménez⁵, Ileana Hernández¹, Dianeys Gonzáles¹, Ramona Márquez¹

¹Departamento de Fisiología y Bioquímica Vegetal
Instituto Nacional de Ciencias Agrícolas, INCA
La Habana, Cuba

²Estación experimental del Arroz
Instituto Nacional de Ciencias Agrícolas
Los Palacios, Pinar del Río, Cuba

³Departamento de Protección de Plantas
Centro Nacional de Sanidad Agropecuaria, CENSA
La Habana, Cuba

⁴Departamento de Biología Celular Vegetal
Facultad de Agricultura, Universidad Notre Damme de la Paix Namur, Bélgica

⁵Facultad de Agronomía
Universidad de Granma, Cuba
E- mail: alfalcon@inca.edu.cu

ABSTRACT

Studies were carried out as part of the Agriculture Biotechnology program, to prepare and characterize chemically and biologically different chitosans obtained from Cuban lobster chitin. Chitosan polymers were subjected to acid and enzymatic hydrolysis by using low-cost commercial enzymatic preparations, and the resulting oligosaccharide mixtures were further characterized. Their potential antimicrobial activities were also evaluated versus fungi and oomycetes, also testing their ability to induce defensive and protective responses in tobacco and rice plants against two economically relevant pests, *Phytophthora nicotianae* and *Pyricularia grisea*, respectively. With the aid of international collaboration, different oligochitosans mixtures were compared for activating defensive responses in suspension cultures of *Arabidopsis thaliana* cells. These results bring knowledge on the physical-chemical properties of the chitosans obtained, such as molar mass and acetylation grade, and their influence on activating defensive responses, the inhibition of growth in pathogens and the induction of resistance in tobacco and rice plants. Some of these chitosan derivatives were selected as possible active components to protect both type of cultivars, being applied at field-scale to evaluate their effects for the main natural pathogens and bringing very promising results. This research allowed us to establish a methodology for preparing oligochitosans, and results shown in here were part of BSc, MSc and PhD theses, and were also published in more than 20 scientific papers and presented in more than 40 scientific conferences.

Keywords: chitosan, enzymatic hydrolysis, antifungal activity, induced resistance, *Phytophthora nicotianae*

Introduction

Higher plants are able to initiate defensive reactions in response to pathogen's infections [1]. Those responses are triggered when the plant recognizes several signals which are released as products of the mutual enzymatic degradation of cell walls of both, the plants and the pathogens [2]. The resulting oligosaccharides, also known as oligosaccharins, not only serve as primary signals of the defensive responses, but also influence other biological responses regarding vegetal growth and development [3, 4]. Among oligosaccharins, the cell wall pectins' oligogalacturonides, β -glucans, and the fungi cell wall chitin and chitosan fragments are potential inducers of resistance and protection from diverse pathogens in plants. That's the reason why they are currently included as active components in several protective products for agriculture [3-6].

Chitin is a linear polymer of N-acetyl-glucosamine, and the second more abundant natural polymer after cellulose. It is found on fungi cell walls and exoskeletons of arthropods as the main production source

worldwide, with more than 10 gigatons (1×10^{13} kg) produced per year. It is mainly obtained from shrimp and crab exoskeletons, with a very small production from lobster exoskeletons [7]. The latter are discarded in large amounts as a byproduct to the sea to avoid environmental problems, which could be used to obtain macromolecules highly applicable in medicine, industry and agriculture [7].

Chitosan, the main chitin derivative, is a linear polymer of glucosamine monomers linked by β 1-4 bonds and it is obtained by alkaline deacetylation. This process produces a polymer soluble in diluted acids, which is advantageous over chitin for agricultural use. On this field, chitosans and their derivatives of lower molecular weights could be applied, based on their proven biological potentialities, such as significant antimicrobial activity on the growth and development of fungi, bacteria and oomycetes [5, 8]; protection of plants from potential pathogens and promotion of cultivar growth and development [5, 6].

1. Agrios GN How plants defend themselves against pathogens (Chapter six). In: Plant Pathology. 5th ed., New York, USA: Academic Press, 2005, p. 208.

2. Esquerré-Tugayé MT, Boudart G, Dumas B. Cell wall degrading enzymes, inhibitory proteins, and oligosaccharides participate in the molecular dialogue between plants and pathogen. *Plant Physiol Biochem* 2000;38:157-63

3. Ridley BL, O'Neill MA, Mohnen D. Pectins: structure, biosynthesis, and oligogalacturonides-related signaling. *Phytochemistry* 2001;57:929-67

4. Shibuya N, Minami E. Oligosaccharide signalling for defenses responses in plant. *Physiol Mol Plant Pathol* 2001;59:223-33

5. Bautista-Baños S, Hernández-Lauzardo AN, Velázquez-del Valle MG, Hernández-López M, Ait Barka E, Bosquez-Molina E, et al. Chitosan as a potential natural compound to control pre and postharvest diseases of horticultural commodities. *Crop Prot* 2006;25:108-18.

✉ Corresponding author

Both, the direct antimicrobial activity and the induction of defensive responses against pathogens of chitosans, are influenced by the physical-chemical properties of the molecule, including the acetylation degree (AD) and molecular weight (MW)[5, 9]. The lower AD of the given chitosan polymer the higher inhibitory activity on microbial growth is obtained [5]. On the contrary, a decrease in the MW of the given chitosan may increase or decrease its inhibitory activity on growth, depending on the specie [5, 10]. These types of studies are scarce on oomycetes, and there are no studies in literature analyzing the influence of these two variables on the development of this genus.

These properties also influence the activation of defensive responses in plants. Kauss and coworkers [11] demonstrated that partial acetylation and fragmentation of chitosans were required for inducing H₂O₂ in cucumber. On the other hand, results from Vander et al. [9] indicated that chitosan polymers with AD over 35% induced the highest Phenylalanine ammonia lyase (PAL) and peroxidase (POD) levels in wheat leaves. Previous works on the role of AD and MW for chitosan-based induction of defensive responses were carried out only in isolated parts of pre- [9] or non-conditioned [11] plants but neither in complete plants nor in the presence of pathogens.

Even when there are reports of the protection of cultures against pathogens by administering different types of chitosans [5, 12], the effect of this compound on the *Nicotiana tabacum-Phytophthora nicotianae* interaction remains to be elucidated, being only reported the partial protection against the Tobacco Mosaic Virus (TMV) by using oligochitosans mixtures [13]. Therefore, there was no information about the induction of resistance against pathogens other than the TMV in tobacco plants, nor in the evaluation of chitosan for inducing responses in Cuban tobacco and rice species.

Given the current context of Cuban agriculture, it is highly relevant to have a non-toxic natural product in origin, produced from national sources and by affordable means, as a replacement alternative for the expensive chemical pesticides being imported.

The Group of Bioactive Products (GPB) at the National Institute of Agricultural Sciences (INCA) has experience in developing methodologies to prepare oligosaccharins from national raw materials and evaluating them as hormone substitutes for *in vitro* culture, to promote root anchoring, soybean symbiosis improvement and induction of plant resistance against diseases. Several of these results have been already patented.

Based on what mentioned above, our research of chitosans focused on the following objectives: development and adaptation of methodologies to obtain chitosans polymers and oligomers from nationally-produced lobster chitin, which show potential biological activity in plants; to evaluate the effect of chitosans of different AD and MW in the *in vitro* development of pathogens of economically relevant crops; to evaluate the effect of chitosans in inducing resistance in suspension cultures of *Arabidopsis* cells, and in tobacco and rice plants inoculated with pathogens and; to test promising compounds on field experiments for both crops.

Materials and methods

Preparation of chitosan poly- and oligosaccharides

Chitosan polymers were obtained by basic deacetylation with NaOH at reactor scale, from nationally obtained lobster chitin. The methodology formerly described to obtain shrimp chitin was adapted for lobster chitin processing at the GBP (Group of bioactive products) of INCA. Studies were carried out by applying chitosan hydrolysis with commercial enzymatic mixes and using typical enzymology techniques, such as: the influence of different substrates at different concentrations, pH, temperature and others. Chitosan mixtures were characterized using mass spectrometry technique (MALDI-TOF) for the first time to determine oligosaccharide AD and MW simultaneously. Acid hydrolysis [14] was also tested to obtain oligochitosans.

Analysis of the effect of chitosans on pathogen growth

The effect of chitosan-based compounds on mycelial growth, sporulation and spores viability were analyzed. Pathogens were isolated from rice and tobacco plants, and cultured in petri dishes filled with solid culture media (PDA, PDA-V8 and/or Czapek Dox culture media) containing the tested chitosan compounds. Spores were incubated with chitosans prior to culturing them on chitosan-free media or by direct germination of spores in chitosan-containing aqueous solutions.

Evaluation of chitosan-induced plant resistance

The activation of defensive responses was studied in suspension cultures of *Arabidopsis thaliana* cells, and in leaves and roots of tobacco and rice plants. PAL induction and the increase of H₂O₂ production in response to chitosan oligosaccharides were determined. According to each experiment, β 1-3 glycanase, chitinase, PAL and POD enzymatic activities were assessed in leaves and roots of tobacco and rice plants. Additionally, the induction of resistance by applying chitosans of different AD and MW was determined in tobacco seedlings against *P. nicotianae* at bioassay scale. It was also tested in rice, by immersing the seeds into a solution containing chitosan polymers and oligomers prior to sowing, and further inoculating the leaves of the resulting seedlings with *Pyricularia grisea*.

Field-scale application of chitosans

Field experiments were carried out in tobacco (variety Habana 92) and rice (variety J-104) cultivars under productive conditions. Tobacco plants were applied with compounds, by foliar aspersion of a chitosan polymer following plant transplantation, and rice cultivars were established by immersing the seeds into a solution containing a chitosan polymer and a chitosan hydrolysate prior to sowing. The natural infection by major pathogens for these two crops was evaluated as indicated on the instructives for infection scales and analyses.

Results and discussion

A methodology similar to that described for obtaining chitosan from shrimp chitin was used to obtain

6. Sharathchandra RG, Niranjan Raj S, Shetty NP, Amruthesh KN, Shetty HS. A Chitosan formulation ElexaTM induces downy mildew disease resistance and growth promotion in pearl millet. *Crop Prot* 2004;23:881-8.

7. Prashanth KVH, Tharanathan RN. Chitin/chitosan: modifications and their unlimited application potential-an overview. *Trends Food Sci Technol* 2007;18:117-31

8. Xu J, Zhao X, Han X, Du Y. Antifungal activity of oligochitosan against *Phytophthora capsici* and other plant pathogenic fungi *in vitro*. *Pest Biochem Physiol* 2007;87:220-8.

9. Vander P, Varum KM, Domard A, El Gueddari NE, Moerschbacher BM. Comparison of the ability of partially N-acetylated chitosans and chitoooligosaccharides to elicit resistance reactions in wheat leaves. *Plant Physiol* 1998;118:1353-9.

10. Jeon Y-J, Shahidi F, Kim SK. Preparation of chitin and chitosan oligomers and their applications in physiological functional foods. *Food Rev Int* 2000;16:159-76.

11. Kauss H, Jeblick W, Domard A, Siegrist J. Partial acetylation of chitosan and a conditioning period are essential for elicitation of H₂O₂ in surface-abraded tissues from various plants. In: Domard A, Roberts GAF, Varum KM (eds), *Advances in Chitin Science II*, Lyon, France.; J André Publisher, 1997, p. 94-101.

12. Trostel-Aziz P, Couderchet M, Vernet G, Aziz A. Chitosan stimulates defense reactions in grapevine leaves and inhibits development of *Botrytis cinerea*. *Eur J Plant Pathol* 2006;114:405-13.

13. Zhao XM, She XP, Du YG, Liang XM. Induction of antifungal resistance and stimulatory effect by oligochitosan in tobacco. *Pest Biochem Physiol* 2007;87:78-84.

chitosan polymers at bank scale. The technique was modified according to a previous study of the influence of chitin/alkali ratio, temperature and reaction time on the properties of the chitosan obtained. Results showed that it is possible to generate high molar mass chitosans under homogeneous preparation conditions (low temperature and an almost equivalent alkali/chitin ratio), with minor modification of its secondary structure and with yields between 74 and 80% of the source chitin. All these are economically relevant for an efficient use in agriculture.

On the other hand, both acidic and enzymatic hydrolyses were studied to generate low MW chitosan compounds. Currently, the enzymatic hydrolysis is preferred over the acidic one, due to its sequence specificity for lysis and the respective formation of products [7]. Nevertheless, the costs of the enzymes used must be low enough to make it affordable for agricultural application and to obtain products with the proper biological activity. The first studies using this strategy showed that it is possible to generate chitosan hydrolyzates having protective activity in plants, by using low cost, commercial enzyme mixes, to improve the biological activity of the obtained chitosan derivatives. In our study, five enzymatic complexes were assayed; allowing us to select the most adequate and it was possible to optimize the hydrolysis conditions (Figure 1). Hydrolyzates and their mixes were obtained with high content of bioactive oligosaccharides and containing the selected complex (Pectinex Ultra SPL) [14].

The biological activity of chitosan oligosaccharides depends on its structure. Otherwise, previous reports

did not address the influence of AD on the activity of quito-oligosaccharides while in others the AD was not determined or ambiguous methods were used to characterize it.

In this work, chitosan was depolymerized by acidic or enzymatic hydrolysis, by using the Pectinex Ultra SPL complex. The resulting oligosaccharides were selectively precipitated in methanol solutions and rigorously characterized by mass spectrometry (MALDI-TOF). The differences in the polymerization degree (PD) and AD were well established (Table 1). The acidic hydrolysis produced fragments with PD up to 16 glucosamine residues, mostly monoacetylated. More significantly, the enzymatic hydrolysis rendered shorter fragments with a high rate of fully deacetylated chitoooligomers, showing a higher oligo size/yield ratio. Based on these evidences, we chose the enzymatic method to prepare oligochitosans [14].

Among the biological assays, the effect of different chitosans on growth and asexual reproduction of to-

14. Cabrera JC, Van Cutsem P. Preparation of chitoooligo-saccharides with degree of polymerization higher than 6 by acid or enzymatic degradation of chitosan. *Biochem Eng J* 2005;25:165-72.

Table 1. Physical-chemical properties of the chitosan polymers and oligomers used in biological assays against *Phytophthora nicotianae*, and the induction of tobacco seedlings resistance against the same pathogen

Type of Chitosan	Abbreviation	Percentage of acetylation degree	Molar mass (g/mol ⁻¹)	Polymerization degree
Polymer	Q-88	12.0 ^a	1.35 x 10 ^{5c}	813 ^d
Polymer	Q-63	36.5 ^a	1.40 x 10 ^{5c}	794 ^d
Oligomers	OLG	0-1.0 ^b	--	5-9 ^b

^aDetermined by infrared spectroscopy

^bDetermined by MALDI-TOF

^cMean molar mass as determined by viscosimetry.

^dAverage degree of polymerization, calculated from the mean molar mass determined by viscosimetry and the monomer molar mass.




Figure 1. Optimization of enzymatic hydrolysis with the enzyme complex Pectinex Ultra SPL. Influence of pH (A), temperature (B), and thermal stability over time (C) of the complex's chitosanase activity on the Q-63 polymer as substrate.

bacco and rice pathogens was evaluated *in vitro*. Significantly novel results were obtained regarding the effect of chitosans at different lifecycle stages of the *Sarocladium* and *Bipolaris* genera. Similarly, it was evidenced the influence of chitosan MW and AD on the different stages of the *P. nicotianae* tobacco pathogen, achieving the highest inhibition by decreasing MW down to oligomers and at the lowest AD. (Figure 2, Table 2). This is a highly significant result, due to its practical relevance when preparing chitosan compounds with inhibitory activity.

It was also evaluated the influence of a chitosan polymer (Q-88) on other pathogens growth (*Phytophthora aphanidermatum*, *Sclerotium rolfsii* and *Rizoctonia solani*) which were isolated from tobacco and other solanaceae, with the first two pathogens as the most sensitive to that polymer, and studying Q-88 for the very first time in dose-response experiments.

Based on these results, and also considering the natural origin and biodegradable nature of chitosan, these polymers and their derivatives were recommended to be used for controlling soil phytopathogens in tobacco seedbeds and to reduce the spotted grain disease in rice.

It was also demonstrated that it is possible to re-acetylate the high PD polymers without affecting their PD, by working with two sets of chitoooligosaccharides of defined PD. Their capacity to induce resistance was evaluated in suspensions of *Arabidopsis thaliana* cells, by measuring the induction of two well known defensive markers: PAL activity and production of hydrogen peroxide (H_2O_2). In this case, the completely deacetylated chitoooligosaccharides induced the activation of both PAL activity and H_2O_2 production, and also cell death, which varied with their PD and concentration. The ability of the oligosaccharides to increase H_2O_2 production and cell death was progressively inhibited by re-acetylation, but the PAL activity was unaffected. This evidenced the role of PD for generating defensive responses on vegetal cells.

Experiments run in tobacco plants showed, for the very first time, the induction of systemic responses against *P. nicotianae* by foliar aspersion, substrata

application and previous treatment of seeds with the chitosan compounds. All these demonstrate the potentiality of these compounds to protect tobacco from its main disease at seedbed level. The induction of resistance, either by activating defensive responses or by reducing the infection of *P. nicotianae* was influenced by the concentration, MW and AD of the applied compounds, and also by the administration procedures [15].

This study evidenced the benefits of protecting tobacco at field scale under semi-controlled conditions, and also the practical relevance of preparing chitosans with the adequate physical-chemical properties, to enhance the protective effect.

In rice, the treatment of seeds of the commercial variety J-104 sensitive to Piriculariosis (*Pyricularia grisea*) with chitosan polymers or hydrolizates prior to sowing, activated defensive responses in plants either or not inoculated with the pathogen, protecting the plants against infection. These assays were carried out under semi-controlled conditions and demonstrated that it is possible to protect this crop from this significant disease at field scale with both types of compounds.

The most promising among all the chitosan-based compounds studied in this work at field scale and under controlled or semi-controlled conditions were assayed at open field scale in both crops. In rice, both the polymer and the hydrolizate were studied, under conditions favoring the occurrence of Piriculariosis, with protection being evidenced only for the chitosan hydrolizate at the concentration assayed.

15. Falcón AB. Chitosanas en la inhibición *in vitro* de *Phytophthora nicotianae* Breda de Haan y en la inducción de resistencia en plantas de tabaco contra este patógeno. Doctoral Thesis in Biological Sciences. Faculty of Biology, University of Havana, July 2009.

Table 2. Effects of chitosans and oligochitosans on *in vitro* viability of *P. nicotianae* zoospores (2.1×10^4 spores/mL)

Chitosan compounds	Compound concentration (g/L)				
	0.5 ^a	1.0	1.5	2.0	2.5
Q-63	+ ^a	+	+	+	-
Q-88	+	+	-	-	-
OLG	+	+	+	-	-

^aThe positive sign indicates mycelia growth in the plate for the seven days next to soaking zoospores with the treatment solutions (soaking for 6 hrs).




Figure 2. Effects of chitosans (Q-63, Q-88) and oligochitosans (OLG) on *in vitro* growth of *P. nicotianae* mycelia (A) in PDA-V8 medium, and of the Q-88 polymer on zoospore viability (1.9×10^4 spores/mL) of *P. nicotianae* at different soaking times (2, 4 and 6 hrs) with the chitosan polymer (B). The result of the bifactorial ANOVA in the anti-mycelial assay, with 8 replicates per treatment, indicated the interaction among the factors at $p \leq 0.05$ (standard error = 0.06). The percentages of inhibition correspond to the original data. Different letters represent significant differences among means of treatments, as determined by comparing them by the Tukey's test ($p \leq 0.05$).

In tobacco, taking together the results of resistance induction and antimicrobial activity, it was decided to evaluate a chitosan polymer (Q-88) by foliar aspersions to plants (variety Habana 92), several days after transplantation and without altering the protection schedule used by the local private producer. In this sense, plants received the chitosan (three doses) and also the scheduled chemical treatment, being finally evaluated for natural infection by viruses, fungi and oomycetes. A significant protection was obtained against all the evaluated pathogens as compared to the control, showing a dose-dependent response for the applied chitosan and also a positive effect of the compound on production yields.

Due to its dual biological action, chitosan-mediated protection of plants can be mediated both by directly affecting pathogen growth and by activating the systemically induced resistance (SIR) in plants. In rice, where the seeds were treated before sowing, the protection found was the second in magnitude. But in tobacco, where the compound was disseminated by aspersions, both effects can be taking place, the antimicrobial action (especially against air-borne pathogens which can get into contact the chitosan adhered to the leaf) and the SIR. The latter evidenced by the reduced infection with root and stem pathogens after aspersions. Both mechanisms of action are vital to implement a protective strategy in crops.

Conclusions

In this work, methodologies were adapted to generate different chitosan compounds from lobster chitin. The resulting and promising polymers, and their mixes and derivatives, were evaluated for biological activity and activation of resistance and protection from pathogens in tobacco and rice plants. It was demonstrated, for the very first time, the influence of certain physical-chemical properties of chitosan derivatives on inhibiting pathogens growth and development and also in the biology of protection of fully grown plants against them. Our results evidenced the structure-

activity relationship among the derivatives prepared and characterized and also the biological functions evaluated.

The first results on protection of rice and tobacco cultivars at field scale are also shown, evidencing their relevance to design and prepare new natural, non-toxic and nationally produced bio-pesticides to be used in agriculture. The inclusion of chitosan as part of the integral management of pathogens would allow reducing or substituting the application of chemicals in some economically relevant crops, with the subsequent protection of the environment.

Scientific relevance of the study

This work provides evidences on the potentialities of chitosan polymers, partially hydrolyzed chitosan and oligochitosans developed in Cuba, to protect tobacco and rice cultivars from relevant diseases which limit production yields. This supports their potential introduction for the ecological control of pathogens in both crops.

By these means, a significant amount of lobster exoskeleton is used as source for chitin and chitosan production, two compounds of high aggregated value, instead of discarding it into the environment with the associated contamination.

Acknowledgements

The authors want to thank the contribution and support of the following collaborators: The technical team at the INCA involved in this work; Dr. Pierre Van Cutsem (Namur University, Belgium), Dr. Silvia Bautista-Baños (IPNY, Morelos, Mexico); Dr. Miguel A Martínez-Téllez (CIAD, Sonora, Mexico); Dr. Fernando Guridi (UNAH); Dr. Eduardo Ortega (University of Havana, Cuba); Dr. Ondina León (CENSA, Cuba); Dr. Verónica Toledo (IIT, Cuba) and Dr. María C. Nápoles (INCA, Cuba). The authors are also grateful to MSc. Regla M Cárdenas and MSc. Elizabeth Cristo; and to the CITMA and MES ministries for partially funding the research included in this work.