

Cuban Journal of
Forest Sciences

CFORES

Volume 11, Issue 2; 2023

*Evaluation of stem profile models of the species *Brachystegia floribunda* (BENTH)*

*Evaluación de modelos de perfil del fuste de *Brachystegia floribunda* (BENTH)*

*Avaliação de modelos de perfil de caule de *Brachystegia floribunda* (BENTH)*

Edilio Aldana Pereira^{1*} , Yordan Lores Pèrez² , Marisela Frías Tamayo¹ 

¹ University of Pinar del Río "Hermanos Saiz Montes de Oca". Pinar del Río, Cuba.

² University of Guantánamo Faculty of Agronomy and Forestry. Guantánamo, Cuba.

*Corresponding author: aldana@upr.edu.cu

Received: 01/30/2023.

Approved: 06/16/2023.

SUMMARY

Brachystegia floribunda (BENTH) has a high economic and social value for the peasant communities of Angola and several African countries. The stem taper or profile functions make it possible to estimate the diameter and/or volume at a given height of the tree, being, therefore, one of the dendrometric characteristics that defines the shape of the stem of forest species. Hence, the present study seeks to evaluate different mathematical models and determine the one with the best fit for the estimation of diameters and volume of this



species. In total, 66 trees were felled and rigorously boxed, of which 53 were used to adjust the stem profile equations and 13 were used to validate the results obtained. Thirteen stem profile models were evaluated and the fit was examined from the comparison of the coefficient of determination, the root mean square error and a precision index. The Clutter non-linear model turned out to be the one with the best precision and, through its mathematical integration, allows obtaining an expression of the taper of the stem compatible with the commercial and total volume of standing trees. This equation is recommended to estimate the diameter and height per section of the stem, as well as the commercial and total volume of the studied species.

Keywords: Stem taper, volumetric estimation, Clutter model, Stem profile, Shape Factor, Shape Quotient, *Brachystegia floribunda*.

RESUMEN

Brachystegia floribunda (BENTH) tiene un alto valor económico y social para las comunidades campesinas de Angola y varios países africanos. Las funciones de ahusamiento o perfil del fuste permiten estimar el diámetro y/o el volumen a una altura determinada del árbol, siendo, por tanto, una de las características dendrométricas que define la forma del fuste de las especies forestales. De ahí es que el presente trabajo persigue evaluar diferentes modelos matemáticos y determinar el de mejor ajuste para la estimación de diámetros y volumen de esta especie. En total, se derribaron y cubicaron rigurosamente 66 árboles, de los cuales fueron utilizados 53 para ajustar las ecuaciones de perfil del fuste y 13 se emplearon para la validación de los resultados obtenidos. Fueron evaluados 13 modelos de perfil del fuste y el ajuste se examinó a partir de la comparación del coeficiente de determinación, la raíz del error medio cuadrático y un índice de precisión. El modelo no lineal de Clutter resultó ser el de mejor precisión y, mediante su integración matemática permite obtener una expresión del ahusamiento del fuste compatible con el volumen comercial y total del arbolado en pie. Esta ecuación es recomendada para estimar el diámetro y altura por sección del fuste, así como, el volumen comercial y total de la especie estudiada.



Palabras clave: Ahusamiento del fuste, estimación volumétrica, modelo de Clutter, perfil del fuste, Factor de Forma, Cociente de Forma, *Brachystegia floribunda*.

RESUMO

A *Brachystegia floribunda* (BENTH) tem um alto valor econômico e social para as comunidades rurais de Angola e de vários países africanos. As funções de taper ou perfil do fuste permitem estimar o diâmetro e/ou o volume a uma determinada altura da árvore, sendo, portanto, uma das características dendrométricas que definem a forma do fuste das espécies florestais. Assim, o presente trabalho tem como objetivo avaliar diferentes modelos matemáticos e determinar o melhor ajuste para a estimativa de diâmetros e volumes dessa espécie. No total, 66 árvores foram abatidas e cubadas rigorosamente, das quais 53 foram utilizadas para ajustar as equações de perfil do fuste e 13 foram utilizadas para validação dos resultados obtidos. Treze modelos de perfil do fuste foram avaliados e o ajuste foi examinado pela comparação do coeficiente de determinação, do erro quadrático médio e de um índice de precisão. O modelo não linear de Clutter foi considerado o mais preciso e, por meio de sua integração matemática, permite obter uma expressão da conicidade do tronco compatível com o volume comercial e o volume total em pé. Essa equação é recomendada para estimar o diâmetro e a altura por seção do caule, bem como o volume comercial e total das espécies estudadas.

Palavras-chave: Conicidade do fuste, estimativa de volume, modelo de Clutter, perfil do fuste, fator de forma, quociente de forma, *Brachystegia floribunda*.

INTRODUCTION

According to Sanfilippo (2014) *Brachystégia floribunda* (BENTH) is a common and exclusive tree species of the “ Miombo ” forest formation of the Central High Plan of Angola. It appears as an arboreal space with a trunk, generally straight and cylindrical, 6-19 cm in normal diameter at 1.3 m from the ground, with rough and irregular bark. According to the information obtained, it is an undemanding species in terms of soil quality that is generally



found as a homogeneous population in soils poor in organic matter, ferralitic, which is characteristic of the region, well-drained and sandy. It produces round wood for rustic constructions and firewood for good quality charcoal. In the forestry aspect, it should be noted that the natural forest mass of Miombo in the village of Calombo where the present investigation was carried out, occupied its entire territorial extension, but gave way, in large part, to savannah plant communities due to human occupation.

The forest mass in the study area is characterized by being made up entirely of natural trees, forming forest patches where the three genera mentioned above predominate, with *B. floribunda* being one of the most representative species. In the Miombo formation of the Angolan Planalto and in the area that is the object of this research, recent works have been carried out and published by Frías Tamayo M. *et al.* (2021), Aldana and Lores (2021), Salvador JA N. *et al.* (2017).

Direct estimation of the volume of individual trees is crucial in forest inventories; for this reason, it is necessary to have mathematical expressions that allow modeling the taper and estimating the total or partial volumes of the stem, through easily measured variables.

According to Pellico (2017) there is extensive information based on the mathematical methodology used to improve the volumetry of trees, always based on the volume of regular solids.

The profile of the stem or taper of a tree is defined as the coefficient of narrowing of the diameter with the increase in height; therefore, the taper or profile functions describe the expected diameters, with or without bark, at different heights in the stem (Ramírez-Martínez A. *et al.*, 2018).

The variation of the taper in the stem of the tree is caused by the natural conditions of growth, and by the response of the forest to the silvicultural treatments applied (Hernández-Pérez *et al.*, 2013 and Uranga-Valencia *et al.*, 2015). The taper can be described by means of a mathematical function that allows predicting the diameters at different heights of the stem; predict the height for a given stem diameter and predict the total volume of the stem or between different stem heights by integrating the model (López *et al.*, 2016). Taper models are an important tool for determining timber stocks (by integrating the taper equation



between two heights) and classifying standing tree products (Ramírez-Martínez A. *et al.*, 2018).

According to Burkhart and Tomé (2012) there are several models that have been used to describe the stem profile, and they have been classified according to the complexity of the mathematical expressions, the number of variables and the coefficients of the model; the general classification is based on: general models for individual tree, segmented models and variable exponent models.

Stem profile analytical curve

The stem profile curves are mathematical functions that allow the average profile of the trees to be represented. The distribution of the diameter along its axis is known as the profile of the trunk or stem and as the curve of the taper of the stem is the curve with which it is intended to represent it; for its part, the directrix curve is the curve generated by a model.

The functions to describe the profile of the stem of the trees was a topic of great interest for foresters (Bi, 2000; Fang *et al.*, 2000; Kozak, 2004). The construction of a volume equation for the classification of different products from a function of the stem profile is based on the capacity of this function to describe the diameter at different heights. Taking into account the application of the definite integral to calculate the volume of a solid of rotation, it is possible to determine the volume of wood between any two heights. Once the limit diameter up to which the volume is to be known is specified, its corresponding height is determined by analytically inverting the function or by means of some iterative procedure (Diéguez-Aranda *et al.*, 2003 and 2006).

The most widely used procedure for estimating individual volume is the use of equations in which volume is the dependent variable, associated with independent variables that are easy to measure in the forest, such as normal diameter at 1.3 m from the ground and height. (Machado, 2002) .

The most common approach on volume compatibles developed systems was carried out to express the β coefficient of Spurr 's combined variable equation, without the independent term ($V = \beta d^2 h$), where V is the total volume of the tree, d is the normal diameter, and h is the total height of the tree) in profile equation form, or vice versa, using a compatibility



relationship. This ensures that the volume function and the profile function are analytically consistent (Sharma and Oderwald , 2011).

There must always be a function that represents the section of the shaft at different heights. Once this function is defined, the total volume of the piece or a part of it can be calculated between two heights h_1 and h_2 . For example, Kozak (2004) used the model (Equation 1) for certain conifers.

$$\frac{d_h^2}{DAP^2} = b_0 + b_1 \left(\frac{h}{H}\right) + b_2 \left(\frac{h}{H}\right)^2 \quad (1)$$

Where: d_h = diameter at height h ;

DAP = diameter at breast height;

h = height at which the diameter of a section is measured;

and H = total height.

Pompas *et al.* (2009) tested six compatible taper models cited in the literature for *Pinus arizonica* engelm , whose expressions correspond to linear and non-linear equations.

Although there is abundant information on the taper of the stem profile of trees, there is little information on equations that describe the stem profile for broadleaf species, whether natural or planted, since most of the stem profile models have been tested on species of the genus *Pinus*.

Despite the efficiency of some equations, they do not always adjust to all forest species and conditions of forest populations, so it is advisable to test them, through statistical trials, and choose the model with the best result (Thomas, 2010).

Most of the methodologies developed for tree volume estimation consider that if the volume of a tree is correctly determined, the found value is valid for another tree of the same diameter, height and shape (Thiersch *et al.*, 2006). The equations of tree taper with height for forest plantations are important to determine the products to be extracted from thinning or other silvicultural interventions. Forest products of different types can be objectively determined with these mathematical technologies. Currently, the functions of diameter taper with height are popular and efficient to represent the stem profile of trees and to estimate, by integration, commercial and total volumes (Clutter *et al.*, 1983 and Nívar, 2013).



There is little research on this subject in the forest species of the Miombo forest formation in the African countries where it exists. Therefore, the scientific problem to be solved in this research is related to the almost complete absence of stem profile for the main Miombo forest species. Hence the need to investigate in this line, with a view to finding models and stem profile equations that allow the precise estimation of the timber existence as support for the sustainable management of Miombo forests. Henceforth, the object of study is the stem profile models applied to the species *B. floribunda* (Benth) in a specific forest area of the Miombo formation.

The objective is to evaluate different mathematical models and determine the one with the best adjustment for the estimation of diameters and volume of *B. floribunda*.

MATERIALS AND METHODS

The present investigation was carried out in the village of Calombo in the Cuima Commune, Caála municipality, Huambo province in Angola, located approximately 25 km from Cuima between the geographical coordinates 13° 30' 12" South latitude and 15° 41' 24" East longitude.

To evaluate the dendrometric and dasometric characteristics, as well as the degree of occupation and productivity of the species *B. floribunda*, 32 circular plots of 500 m² (0.05 hectare) with a radius of 12.62 m were raised. The correction of the radius of the plot was made so that its measurement was adjusted to a flat terrain. For this, the correction was made using the formula $R_i = R_h \cos \alpha$, where: R_i = Radius in the horizontal line; R_h = Radius on the slope line; α = Value of the slope measured in degrees (Figure 1).




Figure 1. - Correction of the radius of the sample plot



For the analysis, a sample of 53 trees was used to fit different stem profile models, and 13 trees were used to validate the best fit model.

Each tree was felled and sectioned, making measurements of the normal diameter ($d_{1.3}$), diameter at the base (db), diameter of the stump at 0.3 m from the ground ($d_{0.3}$), diameters with bark at different heights “taper” (d_{icc}), height (h_i) at different diameters (m) and total height (H) in m. All diameters were measured in cm.

In this study, 1107 pairs of diameters with bark and height data were used, which were obtained from 53 *B. floribunda* trees distributed in a range of diameter and height categories (Table 1).

Table 1 presents the summary of the descriptive statistics of the trunk analyses.

Table 1. - Descriptive statistical values of the studied trees

Variable	No.	Half	Standard deviation	Minimum	Maximum
db	53	12.7	20,545	5	31.9
d_{0.3}	53	11.9	21,176	4.5	30.9
d_{1.3}	53	9.9	14,923	4	18.6
h	53	8.3	2,969	5	13.3
dicc	1107	8.8	19,766	0	23
h_i	1107	3.2	5,451	0	1.0

The dispersion of the heights (hm/H) and diameters (dm/D) in relative terms is illustrated in Figure 2.






Figure 2. - Dispersion of relative diameter and relative height

Tested stem profile models

A stem profile model is a mathematical expression that allows you to predict the diameter of a cross section at any height of the stem and to determine the volume of wood in any segment of the stem. For the adjustment of the stem profile equation, 13 mathematical models were evaluated, of which five are quadratic, five cubic and three non-linear. Eight of the tested models have as dependent variable $\frac{d_{cc}}{d_{1,3}}$, two have $\frac{dh^2}{DAP^2}$ as the dependent variable and in three the dependent variable is d_{cc} (Table 2, appendix 2).

These functions represent the section of the trunk at different heights and once defined allow the calculation of the total volume of the stem or part of it between two heights h_i and h_{i+1} (Equation 2).

Where, in the models, d_{cc} = diameter of the stem with bark at the relative height h (m);

$$h = \text{relative height of the stem above the stump (m)}; X = \frac{(H-h)}{(H-1.30 \text{ m})}; Xk = \frac{(1-Z^{-1/2})}{(1-P^{-1/2})}; \quad (2)$$

$$Z = \frac{h}{H}; H_a = H - 1.30 \text{ m } \beta_1, \beta_2, \beta_3, \beta_4 \text{ are statistical parameters.}$$



For the adjustment of the stem profile functions, 1107 data of diameter and height corresponding to 53 trees were used. For the equations that required transformations, the statistical parameters were calculated with the necessary transformations.

A correlation analysis for the stem profile made it possible to determine the most correlated dependent and independent variables in the stem profile model.

To determine the validation statistics the determinant (R^2) coefficient, the standard error of the estimation (EEE) and the average bias were mainly used, which were calculated as follows Equation 3):

$$Sesgo\ medio = \frac{\sum_{i=1}^n (Y_i - \tilde{Y}_i)}{n}; R^2 = 1 - \frac{\sum_{i=1}^n (Y_i - \tilde{Y}_i)^2}{\sum_{i=1}^n (Y_i - \bar{Y}_i)^2}; EEE = \left[\frac{\sum_{i=1}^n (Y_i - \tilde{Y}_i)^2}{n-p} \right]^{\frac{1}{2}} \quad (3)$$

Where:

Y_i = observed value or the dependent variable;

\bar{Y}_i = mean value of the observed data;

\tilde{Y}_i = predictive value; and

p = number of model parameters including the intercept.

RESULTS AND DISCUSSION

The Microsoft Excel 2010 allowed calculating the main dendrometric and dasometric variables of the species *B. floribunda*. Among these variables are: number of trees per hectare (Nha^{-1}); average diameter at 1.3 m from the ground ($\bar{d}_{1.3}$); diameter at 0.3 m from the ground ($\bar{d}_{0.3}$); mean total height (\bar{h}_t); average commercial height (\bar{h}_c); mean basal area in m^2 per hectare at 1.3 m above the ground ($\bar{G}_{1.3}ha^{-1}$); mean basal area in m^2 per hectare at 0.3 m from the ground ($\bar{G}_{0.3}ha^{-1}$); the average volume in m^3 per hectare with the basal area at 1.3 m from the soil ($\bar{V}_{1.3}ha^{-1}$) and the average volume in m^3 per hectare with the basal area at 0.3 m from the soil ($\bar{V}_{0.3}ha^{-1}$). These results were obtained from the 32 plots surveyed with the aim of characterizing the forest mass where the research was carried out and are



summarized in Table 2, both for the total per hectare for the forest mass and for the respective diameter classes (Table 2).

Table 2. - Main dasometric indices of the species *B. Floribunda*

CD	Nha ⁻¹	$\bar{d}_{1.3}$ (cm)	$\bar{d}_{0.3}$ (cm)	\bar{h}_t (m)	$\bar{G}_{1.3}$ (m ² ha ⁻¹)	$\bar{G}_{0.3}$ (m ² ha ⁻¹)	$\bar{V}_{1.3}$ (m ³ ha ⁻¹)	$\bar{V}_{0.3}$ (m ³ ha ⁻¹)
2	83	23	3.4	2.4	0.001	0.003	0.0029	0.006
4	168	3.9	5.1	3.5	0.021	0.017	0.056	0.042
6	182	5.7	7.9	5.1	0.027	0.045	0.086	0.133
8	201	7.7	9.6	6.0	0.046	0.070	0.163	0.229
10	133	9.8	12.1	6.7	0.049	0.078	0.195	0.290
12	82	11.7	13.5	7.5	0.043	0.056	0.176	0.207
14	55	13.5	15.9	8.1	0.039	0.053	0.167	0.190
16	33	15.1	18.0	9.0	0.029	0.043	0.130	0.149
18	42	17.5	18.8	9.9	0.048	0.059	0.242	0.269
twenty	twenty	19.0	19.5	8.0	0.028	0.030	0.240	0.244
22	twenty	21.0	22.0	10.0	0.035	0.038	0.176	0.089
24	twenty	24.2	22.0	9.0	0.046	0.038	0.215	0.178
30	twenty	29.6	30.9	12.8	0.069	0.075	0.405	0.441
Total/ hectare	1060	13.9	15.3	7.5	9,655	12,112	45,065	49,333

In general, in the area under study, the number of trees per hectare of diameter classes greater than 15 cm is very low.

Occupancy degree by diameter classes

In Table 3, it can be seen that the degree of occupation of *B. floribunda* (Benth) is high, since it occupies 42.9 % of the total number of trees per hectare of the forest mass, and also the average values of $\bar{d}_{1.3}$, $\bar{d}_{0.3}$, and \bar{h}_t are respectively 1.6; 1.5 and 1.4 times higher than the average values of all the species inventoried in the area. 61.5 % and 52.2 % of the average basal area per hectare respectively at 1.3 m and 0.3 m from the soil, as well as 67 % and 49.5



% of the average volume per hectare also at 1.3 m and 0.3 m from the soil respectively corresponds to *B. floribunda* (Table 3).

Table 3. - Degree of occupation of *B. floribunda* in the forest area under study

Parcel	No ha ⁻¹	$\bar{d}_{1.3}$ (cm)	$\bar{d}_{0.3}$ (cm)	\bar{h}_t (m)	$\bar{G}_{1.3}$ (m ² ha ⁻¹)	$\bar{G}_{0.3}$ (m ² ha ⁻¹)	$\bar{V}_{1.3}$ (m ³ ha ⁻¹)	$\bar{V}_{0.3}$ (m ³ ha ⁻¹)
Average values per hectare of all species.	2470	8.8	10.4	5.5	15.7	23.2	67.3	99.7
Average values per hectare from <i>B. floribunda</i> (Benth).	1060	13.9	15.3	7.5	9,655	12,112	45,065	49,333
%	42.9				61.5	52.2	67.0	49.5

This species has a high energy value, which is why it is widely used in the production of charcoal and, according to previous data, it represents a high potential in terms of stocks in irregular Miombo forests for the production of charcoal.

Table 4 shows the percentage of trees in the respective diameter classes, where the highest percentages are observed between classes 4 and 10 with emphasis on diameter classes 8, 6 and 4, in that order respectively (Table 4).

Table 4. - Percentage of tree occupation by diameter classes of *B. floribunda*

diameter classes	Nha ⁻¹	%	diameter classes	Nha ⁻¹	%
2	83	8	16	33	3
4	168	16	18	42	4
6	182	17	20	20	2
8	201	19	22	20	2
10	133	13	24	20	2
12	82	8	30	20	2
14	55	5			
Total/ hectare				1060	100



The diameter distribution of the number of trees per hectare presents the typical shape of irregular or multi-age forests, that is to say, that it has a negative exponential trend of inverted J, which means that the smaller diameter classes occur with a higher frequency of individuals than the smaller diameter superior classes, this distribution guarantees the perpetuity of this natural forest (Figure 3).

This feature is common in most tropical forest formations and has been confirmed by Aldana (2010), but it had previously been explained by Scheffer *et al.* (1930) and Machado (2002).




Figure 3. - Distribution of *B. floribunda* trees by diameter classes

Evaluation of the best profile model of the stem of *B. floribunda*

A correlation analysis of the diameter with bark (d_{cc}) was carried out, measured in different sections along the stem, with respect to the other variables, in order to determine the dependent variable of the general model of the stem profile. The correlation matrix of this.

Analysis is shown in Table 6. This Pearson bivariate correlation analysis allowed us to determine the variables most correlated with the diameter with bark (Table 5).



Table 5. - Pearson's bivariate correlation matrix of the dendrometric variables for the profile of the stem of *B. floribunda*

		db	d _{0.3}	d _{1.3}	h	dcc	hi
d _{cc} ***	Pearson correlation	.724 **	,740 **	,777 **	.434 **	1	-.538 **
	Next (bilateral)	,000	,000	,000	,000	,000	,000
	No.	1112	1112	1112	1112	1112	1112

*** Diameter with bark
 ** Correlation is significant at the 0.01 level (bilateral).
 * The correlation is significant at the 0.05 level (bilateral).

As observed in Table 5, Pearson's variables with the highest correlation are d_{cc} with itself and with $d_{1.3}$. Therefore, the two most correlated variables were chosen, that is, the dependent variables of the model are d_{cc} and $\frac{d_{cc}}{d_{1.3}}$

Determination of the predictive capacity and goodness of fit of the different models

With the application of SPSS Statists version 19, the statistical analysis was carried out, where the parameters, the correlations of the parameter estimates, the ANOVA and the descriptive statistics were estimated, which are basic for the determination of the goodness of fit and the prediction capacity of each model.

In Table 6, (see appendix 2) are the equations resulting from each stem profile model with the main statistical indices. In this table, in addition to the equations with the respective values of the regression coefficients, the following also appear: the correlation coefficient (R), the determination coefficient (R²), the standard deviation (S_x), the root of the error means squares and the aggregate difference. These statistical indices constitute the tool to decide, according to their behavior, which of the tested models will have the best goodness of fit and predictive capacity. Table 6 shows the equations of the five models with the best goodness of fit and predictive capacity.

For the evaluation of each of the five selected models, the tree with the largest number of sections was chosen from the database of the 13 trees defined for validation, and then the diameter of each section was estimated. The calculation of the three statistical indices; Mean bias, Coefficient of determination (R²) and Standard Error of Estimates (SEE), allowed to



carry out the validation and determine the best model of the profile of the stem of the species *B. floribunda*, whose formulas are (Equation 4):

$$Sesgo\ médio = \frac{\sum_{i=1}^n (Y_i - \hat{Y}_i)}{n} ; \quad r^2 = 1 - \frac{\sum_{i=1}^n (Y_i - \hat{Y}_i)^2}{\sum_{i=1}^n (Y_i - \bar{Y})^2} ; \quad EPE = \left[\frac{\sum_{i=1}^n (Y_i - \hat{Y}_i)^2}{n - p} \right]^{\frac{1}{2}} \quad (4)$$

Table 6 shows the statistical validation results of the five models, where it can be seen that model 12, represented by equation

$d_{cc} = 0,069 * d_{1,3}^{1,745} + H^{-0,434}(H - h_i)^{1,391}$, is the best, since it presents the lowest average bias and the lowest standard error of the estimation, in addition to present the highest coefficient of determination (Table 6 and Table 7).

Table 6. - The best 5 models of stem profile with bark tested on the species *B. floribunda*

No	Model Types	R.	R2 -	sx	REC M	GIVE S	equations
2	Cubic	0.868	0.753	0.1348	0.004	0.018	$\frac{d_{cc}}{d_{1,3}} = 1,161 + (-0,558)\left(\frac{h}{H}\right) + (-1,602)\left(\frac{h}{H}\right)^2 + 1,247\left(\frac{h}{H}\right)^3$
4	Cubic	0.860	0.740	0.1390	0.0042	0.019	$\frac{d_{cc}}{d_{1,3}} = 1,024 + (-0,827)\left(\frac{h-1,30}{H-1,30}\right) + (-0,640)\left(\frac{h-1,30}{H-1,30}\right)^2 + 0,694\left(\frac{h-1,30}{H-1,30}\right)^3$
eleven	Quadratic (Amidón)	0.917	0.840	1.7416	.05227	3,033	$d_{cc} = 1,006 * \frac{d_{1,3}(H-h)}{H-1,3} + (-0,118) * \frac{(H^2 - h^2)(h_i - 1,3)}{H^2}$
12	nonlinear (Clutter 1983)	0.919	0.845	1.7454	0.0353	3,046	$d_{cc} = 0,069 * d_{1,3}^{1,745} + H^{-0,434}(H - h_i)^{1,391}$
13	non linear	0.965	0.932	1.1586	0.0349	1,342	$d_{cc} = (0,423 + 1,081d_{1,3}) \left[\frac{H-h}{H-1,3} \right]^{0,956}$

Table 7. - Validation statistics of the 5 best stem profile models of *B. floribunda*

Statistical indices	model 2	model 4	model 11	model 12	model 13
mean bias	0.152	0.173	0.505	-0.099	-0.535
R2 -	0.803	0.804	0.707	0.839	0.710
EEA	0.872	0.869	1,031	0.787	1,041



The graphical representation of the stem profile with each of the five models (Figure 4) shows that models 12 and 13 are the closest to the real value. Therefore, it can be concluded that the equation $d_{cc} = 0,069 * d_{1,3}^{1,745} + H^{-0,434}(H - h_i)^{1,391}$ is the one that best defines the stem profile and most accurately estimates the diameters of the different sections of a *B. floribunda* tree.




Figure 4. - Represents a graph of the stem profile with different equations

CONCLUSIONS

B. floribunda is the most represented species in the study area with 42.9% of the total number of trees per hectare, 61.5% and 52.2 % of the basal area per hectare at 1.3 and 0.3 m from the soil respectively, as well as 67.0 and 49.5 % of the average volume per hectare at 1.3 and 0.3 m from the soil respectively.

In the same way, the mean diameters at 1.3 and 0.3 m from the ground and mean total height are 1.6; 1.5 and 1.4 times higher than the average values of all the species inventoried in the area.

Of the 13 adjusted stem profile models, the non-linear model by Clutter (1983) was the best ,whose general formula is $d_{cc} = b_1 * D^{b_2} + H^{b_3}(H - h)^{b_4}$ by the equation that defines the stem profile of the species *B. floribunda* is (Equation 5):



$$d_{cc} = 0,069 * d_{1,3}^{1,745} + H^{-0,434}(H - h_i)^{1,391}. \quad (5)$$

Appendix 1 (Table 8)

Table 8. - Models adjusted for the profile of the stem of *B. floribunda*

No	Model Types	equations
1	nonlinear quadratic	$\frac{d_{cc}}{d_{1,3}} = b_0 + b_1 \left(\frac{h_i}{h_t} \right) + b_2 \left(\frac{h_i}{h_t} \right)^2$
2	nonlinear cubic	$\frac{d_{cc}}{d_{1,3}} = b_0 + b_1 \left(\frac{h_i}{h_t} \right) + b_2 \left(\frac{h_i}{h_t} \right)^2 + b_3 \left(\frac{h_i}{h_t} \right)^3$
3	nonlinear quadratic	$\frac{d_{cc}}{d_{1,3}} = b_0 + b_1 \left(\frac{h_i - 1.30}{h_t - 1.30} \right) + b_2 \left(\frac{h_i - 1.30}{h_t - 1.30} \right)^2$
4	nonlinear cubic	$\frac{d_{cc}}{d_{1,3}} = b_0 + b_1 \left(\frac{h_i - 1.30}{h_t - 1.30} \right) + b_2 \left(\frac{h_i - 1.30}{h_t - 1.30} \right)^2 + b_3 \left(\frac{h_i - 1.30}{h_t - 1.30} \right)^3$
5	nonlinear quadratic	$\frac{d_{cc}}{d_{1,3}} = b_0 + b_1 \left(\frac{h_i}{h_t} - 1 \right) + b_2 \left(\frac{h_i}{h_t} - 1 \right)^2$
6	nonlinear cubic	$\frac{d_{cc}}{d_{1,3}} = b_0 + b_1 \left(\frac{h_i}{h_t} - 1 \right) + b_2 \left(\frac{h_i}{h_t} - 1 \right)^2 + b_3 \left(\frac{h_i}{h_t} - 1 \right)^3$
7	nonlinear quadratic	$\frac{d_{cc}}{d_{1,3}} = b_0 + b_1 \left(\frac{h_t - h_i}{h_t - 1.30} \right) + b_2 \left(\frac{h_t - h_i}{h_t - 1.30} \right)^2$
8	nonlinear cubic	$\frac{d_{cc}}{d_{1,3}} = b_0 + b_1 \left(\frac{h_t - h_i}{h_t - 1.30} \right) + b_2 \left(\frac{h_t - h_i}{h_t - 1.30} \right)^2 + b_3 \left(\frac{h_t - h_i}{h_t - 1.30} \right)^3$
9	Quadratic (Kozak)	$\frac{dh^2}{DAP^2} = b_0 + b_1 \left(\frac{h}{H} \right) + b_2 \left(\frac{h}{H} \right)^2$
10	Cubic (Kozak)	$\frac{dh^2}{DAP^2} = b_0 + b_1 \left(\frac{h}{H} \right) + b_2 \left(\frac{h}{H} \right)^2 + b_3 \left(\frac{h}{H} \right)^3$
eleven	nonlinear (Amidon 1984)	$d_{cc} = \beta_0 \frac{D(H-h)}{H-1.3} + \beta_1 \frac{(H^2-h^2)(h-1.3)}{H^2}$
12	nonlinear (Clutter 1983)	$d_{cc} = \beta_1 * D^{\beta_2} + H^{\beta_3}(H - h)^{\beta_4}$
13	nonlinear (Rustagi and Loveles 1991)	$d_{cc} = (\beta_0 + \beta_1 d_{1,3}) \left[\frac{H-h}{H-1,37} \right]^{\beta_2}$



Appendix 2 (Table 9)

Table 9. - Models evaluated in the species *B. floribunda* to select the one with the best goodness of fit and predictive capacity

No	Model Types	R.	R2	sx	REC M	GIV ES	Equations
1	Quadra tic	0.8 53	0.7 28	0.14 14	0.004 2	0.020	$\frac{d_{cc}}{d_{1,3}} = 1,225 + (-1,376)\left(\frac{h}{H}\right) + 0,465\left(\frac{h}{H}\right)^2$
2	Cubic	0.8 68	0.7 53	0.13 48	0.004 0	0.018	$\frac{d_{cc}}{d_{1,3}} = 1,161 + (-0,558)\left(\frac{h}{H}\right) + (-1,602)\left(\frac{h}{H}\right)^2 + 1,247\left(\frac{h}{H}\right)^3$
3	Quadra tic	0.8 45	0.7 13	0.14 58	0.004 4	0.021	$\frac{d_{cc}}{d_{1,3}} = 1,001 + (-0,985)\left(\frac{h-1.30}{H-1.30}\right) + 0,298\left(\frac{h-1.30}{H-1.30}\right)^2$
4	Cubic	0.8 60	0.7 40	0.13 90	0.004 2	0.019	$\frac{d_{cc}}{d_{1,3}} = 1,024 + (-0,827)\left(\frac{h-1.30}{H-1.30}\right) + (-0,640)\left(\frac{h-1.30}{H-1.30}\right)^2 + 0,694\left(\frac{h-1.30}{H-1.30}\right)^3$
5	Quadra tic	0.8 52	0.7 25	0.14 21	0.004 3	0.020	$\frac{d_{cc}}{d_{13}} = 1,223 + (-1,181)\left(\frac{h}{H} - 1\right) + 0,341\left(\frac{h}{H} - 1\right)^2$
6	Cubic	0.8 65	0.7 49	0.13 58	0.004 1	0.018	$\frac{d_{icc}}{d_{1,30}} = 1,160 + (-0,484)\left(\frac{h}{H} - 1\right) + (-1,183)\left(\frac{h}{H} - 1\right)^2 + 0,795\left(\frac{h}{H} - 1\right)^3$
7	Quadra tic	0.8 54	0.7 30	0.14 09	0.004 2	0.020	$\frac{d_{cc}}{d_{13}} = 0,314 + 0,377\left(\frac{H-h}{H-1.30}\right) + 3,12\left(\frac{H-h}{H-1.30}\right)^2$
8	Cubic	0.8 68	0.7 54	0.13 44	0.004 0	0.018	$\frac{d_{cc}}{d_{13}} = 0,253 + 0,029\left(\frac{H-h}{H-1.30}\right) + 1,409\left(\frac{H-h}{H-1.30}\right)^2 + (-0,666)\left(\frac{H-h}{H-1.30}\right)^3$
9	Quadra tic (Kozak)	0.6 76	0.4 57	0.47 79	0.014 4	0.228	$\frac{dh^2}{DAP^2} = -0,349 + 4,695\left(\frac{h}{H}\right) + (-3,660)\left(\frac{h}{H}\right)^2$
10	Cubic (Kozak)	0.7 46	0.5 56	0.43 21	0.013 0	0.187	$\frac{d_{h_i}}{DAP^2} = 0,023 + (-0,764)\left(\frac{h}{H}\right) + 11,981\left(\frac{h}{H}\right)^2 + (-10,735)\left(\frac{h}{H}\right)^3$
elev en	Quadra tic (Amid on)	0.9 17	0.8 40	1.74 16	.0522 7	3,033	$d_{cc} = 1,006 * \frac{d_{1,3}(H-h)}{H-1.3} + (-0,118) * \frac{(H^2-h^2)(h_i-1.3)}{H^2}$
12	Nonlin ear (Clutte r 1983)	0.9 19	0.8 45	1.74 54	0.035 3	3,046	$d_{cc} = 0,069 * d_{1,3}^{1,745} + H^{-0,434}(H-h_i)^{1,391}$



13 nonlinear 0.9 0.9 1.15 0.034 1,342
 ar 65 32 86 9

$$d_{cc} = (0,423 + 1,081d_{1,3}) \left[\frac{H-h}{H-1,3} \right]^{0,956}$$

Appendix 5 (Table 10)

Table 10. - Real diameter ($d_{i\ cc}$) and calculated (d_{cc}) with the best five stem profile models

say _{cc}	hi	dcc_ Model 2	dcc_ Model 4	dcc_ Model 11	dcc_ Model 12	dcc_ Model 13
10.8	0	8.9	8.9	9.4	9.6	10.3
9	0.3	8.8	8.7	9.1	9.2	10.0
8.2	0.5	8.6	8.6	8.8	8.9	9.7
8.2	0.5	8.6	8.6	8.8	8.9	9.7
8	0.75	8.4	8.4	8.5	8.6	9.4
8	1	8.2	8.2	8.1	8.3	9.1
8	1	8.2	8.2	8.1	8.3	9.1
7.7	1.25	8.0	7.9	7.8	8.0	8.8
7.7	1.5	7.7	7.7	7.5	7.7	8.5
7.7	1.5	7.7	7.7	7.5	7.7	8.5
6.7	1.75	7.5	7.4	7.2	7.4	8.2
6.6	2	7.2	7.1	6.8	7.2	7.9
6.6	2	7.2	7.1	6.8	7.2	7.9
6.5	2.25	6.9	6.9	6.5	6.9	7.6
6	2.5	6.6	6.6	6.2	6.6	7.3
6	2.5	6.6	6.6	6.2	6.6	7.3
6	2.75	6.3	6.3	5.9	6.3	7.0
6	3	6.0	6.0	5.5	6.1	6.7
6	3	6.0	6.0	5.5	6.1	6.7
6	3.25	5.7	5.7	5.2	5.8	6.3
5.6	3.5	5.4	5.4	4.9	5.5	6.0
5.6	3.5	5.4	5.4	4.9	5.5	6.0
5.6	3.75	5.1	5.1	4.6	5.3	5.7
5.6	4	4.8	4.8	4.3	5.1	5.4
5.6	4	4.8	4.8	4.3	5.1	5.4
5.2	4.25	4.5	4.5	4.0	4.8	5.1
5.2	4.5	4.2	4.2	3.7	4.6	4.8
5.2	4.5	4.2	4.2	3.7	4.6	4.8
4.9	5	3.6	3.6	3.1	4.1	4.1
4.9	5	3.6	3.6	3.1	4.1	4.1
4.9	5.25	3.4	3.4	2.8	3.9	3.8
4	5.5	3.1	3.2	2.5	3.7	3.5
4	5.5	3.1	3.2	2.5	3.7	3.5
3.9	5.75	2.9	2.9	2.2	3.5	3.2
3.9	6	2.7	2.7	1.9	3.4	2.9
6.2		5.9	5.9	5.6	6.2	6.6



BIBLIOGRAPHIC REFERENCES

- ALDANA PEREIRA, E., 2010. *Medición forestal* [en línea]. La Habana: Félix Varela. ISBN 978-959-07-1321-7. Disponible en: https://books.google.com.cu/books/about/Medici%C3%B3n_forestal.html?id=Pv6OswEACAAJ&re%20dir_esc=y.
- ALDANA PEREIRA, E. y LORES PÉREZ, Y., 2021. Factor y cociente de forma preliminar de *Brachystegia floribunda* Benth. *Revista Cubana de Ciencias Forestales* [en línea], vol. 9, no. 1, pp. 124-139. [Consulta: 6 marzo 2023]. ISSN 2310-3469. Disponible en: <https://cfores.upr.edu.cu/index.php/cfores/article/view/637>.
- ÁLVAREZ GONZÁLEZ, J.G., ÁLVAREZ TABOADA, M.F., BARRIO ANTA, M., CASTEDO DORADO, F., DIÉGUEZ ARANDA, U., ROJO ALBORECA, A. y RUIZ GONZÁLEZ, A.D., 2003. *Dendrometría* [en línea]. España: Ediciones Mundi-Prensa. ISBN 978-84-8476-145-7. Disponible en: https://books.google.com.cu/books?id=25dwscrGdeMC&printsec=frontcover&hl=es&source=gbg_summary_r&cad=0#v=onepage&q&f=false.
- AMARAL MACHADO, S. do, BARBOSA DA CONCEIÇÃO, M. y FIGUEIREDO, D.J. de, 2002. Modelagem do volume individual para diferentes idades e regimes de desbaste em plantações de *Pinus oocarpa*. *RECEN-Revista Ciências Exatas e Naturais* [en línea], vol. 4, no. 2, pp. 185-197. [Consulta: 6 marzo 2023]. ISSN 2175-5620. Disponible en: <https://revistas.unicentro.br/index.php/RECEN/article/view/462>.
- BURKHART, H. y TOMÉ, M., 2012. *Modeling forest trees and stands* [en línea]. Estados Unidos: Springer. ISBN 978-94-007-1597-4. Disponible en: https://www.researchgate.net/publication/235004853_Modeling_Forest_Trees_and_Stands.



- CLUTTER, J.L., 1983. *Timber Management: A Quantitative Approach* [en línea]. Estados Unidos: Wiley. ISBN 978-0-471-02961-8. Disponible en: https://books.google.com.cu/books/about/Timber_Management.html?id=RSbxAAAAMAAJ&redir_esc=y.
- DIÉGUEZ ARANDA, U., CASTEDO DORADO, F., ÁLVAREZ GONZÁLEZ, J. y ROJO ALBORECA, A., 2006. Compatible taper function for Scots pine plantations in northwestern Spain. *Canadian Journal of Forest Research* [en línea], vol. 36, no. 5, pp. 1190-1205. DOI 10.1139/x06-008. Disponible en: https://www.researchgate.net/publication/237866454-Compatible_taper_function_for_Scots_pine_plantations_in_northwestern_Spain.
- FANG, BORDERS, B. y BAILEY, L., 2000. Compatible Volume-Taper Models for Loblolly and Slash Pine Based on a System with Segmented-Stem Form Factors. *Forest Science* [en línea], vol. 46, no. 1, pp. 1-12. Disponible en: https://www.researchgate.net/publication/233512687-Compatible_Volume-Taper_Models_for_Loblolly_and_Slash_Pine_Based_on_a_System_with_Segmented-Stem_Form_Factors.
- FRÍAS TAMAYO, M., PEREIRA JAY, N.F. y ALDANA PEREIRA, E., 2021. Evaluación preliminar de la eficacia del método con probabilidad variable en bosque de Miombo. *Revista Cubana de Ciencias Forestales* [en línea], vol. 9, no. 2, pp. 193-204. [Consulta: 6 marzo 2023]. ISSN 2310-3469. Disponible en: <https://cfores.upr.edu.cu/index.php/cfores/article/view/635>.
- HERNÁNDEZ PÉREZ, D., SANTOS POSADAS, H.M. de los, ÁNGELES PÉREZ, G., VALDEZ LAZALDE, J.R. y VOLKE HALLER, V.H., 2013. Funciones de ahusamiento y volumen comercial para *Pinus patula* Schltdl. et Cham. en Zacualtipán, Hidalgo. *Revista Mexicana de Ciencias Forestales* [en línea], vol. 4, no. 16, pp. 34-45. [Consulta: 6 marzo 2023]. Disponible en: <https://www.redalyc.org/articulo.oa?id=63433990004&idioma=en>.



- HUIQUAN BI, 2000. Trigonometric variable-form taper equations for Australian eucalypts. *Forest science* [en línea], vol. 46, no. 3, pp. 397-409. ISSN 0015-749X. Disponible en: <https://pascal-francis.inist.fr/vibad/index.php?action=getRecordDetail&idt=1531809>.
- KOZAK, A., 2004. My last words on taper equations. *The Forestry Chronicle* [en línea], vol. 80, no. 4, pp. 507-515. DOI 10.5558/tfc80507-4. Disponible en: https://www.researchgate.net/publication/272900236_My_last_words_on_taper_equations.
- LÓPEZ AGUIRRE, A., BARRIOS, A. y TRINCADO, G., 2016. Modelos de perfiles fustales con una estructura de error autorregresiva para plantaciones de *Eucalyptus tereticornis* en Colombia. *Madera y Bosques* [en línea], vol. 21, no. 2. DOI 10.21829/myb.2015.212446. Disponible en: https://www.researchgate.net/publication/309138157_Modelos_de_perfiles_fustales_con_una_estructura_de_error_autorregresiva_para_plantaciones_de_Eucalyptus_tereticornis_en_Colombia.
- NANVONAMUQUITXO, S.J.A., GÓNGORA ROJAS, F. y HOFIÇO, N.D.S.A., 2017. Estructura y composición florística de un bosque nativo de Miombo en el Distrito de Mocuba Provincia Zambezia, Mozambique. *Revista Cubana de Ciencias Forestales* [en línea], vol. 5, no. 2, pp. 140-152. [Consulta: 6 marzo 2023]. ISSN 2310-3469. Disponible en: <https://cfores.upr.edu.cu/index.php/cfores/article/view/229>.
- NÁVAR, J., RODRÍGUEZ FLORES, F. de J. y DOMÍNGUEZ CALLEROS, P., 2013. Taper functions and merchantable timber for temperate forests of northern Mexico. *Annals of Forest Research* [en línea], vol. 56, no. 1, pp. 165-178. DOI 10.15287/afr.2013.51. Disponible en: https://www.researchgate.net/publication/286642601_Taper_functions_and_merchantable_timber_for_temperate_forests_of_northern_Mexico.



- OLIVEIRA, A.D. de, DEHON, G., SCOLFORO, J.R., THIERSCH, C. y MAESTRI, R., 2006. Acurácia dos métodos para estimativa do volume comercial de clones de *Eucalyptus* sp. *Cerne* [en línea], vol. 12, no. 2, pp. 167-181. Disponible en: https://www.researchgate.net/publication/237039557_Acuracia_dos_metodos_para_estimativa_do_volume_comercial_de_clones_de_Eucalyptus_sp.
- PÉLLICO NETTO, S., 2017. Equivalência volumétrica: uma nova metodologia para estimativa do volume de árvores. *Revista Acadêmica: Ciência Animal* [en línea], vol. 2, no. 1, pp. 17. DOI 10.7213/cienciaanimal.v2i1.15003. Disponible en: https://www.researchgate.net/publication/321284406_Equivalencia_volumetrica_uma_nova_metodologia_para_estimativa_do_volume_de_arvores.
- POMPA GARCÍA, M., CORRAL RIVAS, J.J., DÍAZ VÁSQUEZ, M. y MARTÍNEZ SALVADOR, M., 2009. Función de ahusamiento y volumen compatible para *Pinus arizonica* Engelm. En el Suroeste de Chihuahua. *Ciencia forestal en México* [en línea], vol. 34, no. 105, pp. 117-134. Disponible en: https://www.researchgate.net/publication/262448053_Funcion_de_ahusamiento_y_volumen_compatible_para_Pinus_arizonica_Engelm_En_el_Suroeste_de_Chihuahua.
- RAMÍREZ MARTÍNEZ, A., SANTIAGO GARCÍA, W., QUIÑONEZ BARRAZA, G., RUIZ AQUINO, F. y ANTÚNEZ, P., 2018. Modelación del perfil fustal y volumen total para *Pinus ayacahuite* Ehren. *Madera y Bosques* [en línea], vol. 24, no. 2. DOI 10.21829/myb.2018.2421496. Disponible en: https://www.researchgate.net/publication/325461053_Modelacion_del_perfil_fustal_y_volumen_total_para_Pinus_ayacahuite_Ehren.
- SANFILIPPO, M., 2014. *Trinta árvores e arbustos do miombo Angolano - Guia de campo para a identificação* [en línea]. Italia: COSPE. [Consulta: 6 marzo 2023]. Disponible en: <http://www.bibliotecaflorestal.ufv.br/handle/123456789/14828>.



SCHEFFER, A., GAZIN, A. y ALVERNY, A. d', 1930. *Sapinières*. Paris: Lwes Presses Universitieres de France.

SHARMA, M. y ODERWALD, R., 2011. Dimensionally compatible volume and taper equations. *Canadian Journal of Forest Research* [en línea], vol. 31, no. 5, pp. 797-803. DOI 10.1139/x01-005. Disponible en: https://www.researchgate.net/publication/237865757_Dimensionally_compatible_volume_and_taper_equations.

THOMAS, C., ANDRADE, C., SCHNEIDER, P. y FINGER, C., 2010. Comparação de equações volumétricas ajustadas com dados de cubagem e análise de tronco. *Ciencia Florestal* [en línea], vol. 16, no. 3. DOI 10.5902/198050981911. Disponible en: https://www.researchgate.net/publication/27790615_Comparacao_de_equacoes_volumetricas_ajustadas_com_dados_de_cubagem_e_analise_de_tronco.

URANGA VALENCIA, L.P., DE LOS SANTOS POSADAS, H.M., VALDEZ LAZALDE, J.R., LÓPEZ UPTON, J. y NAVARRO GARZA, H., 2015. Volumen total y ahusamiento para *Pinus patula* Schiede ex Schltdl. et Cham. en tres condiciones de bosque. *Agrociencia* [en línea], vol. 49, no. 7, pp. 787-801. [Consulta: 6 marzo 2023]. ISSN 1405-3195. Disponible en: <https://dialnet.unirioja.es/servlet/articulo?codigo=5425690>.

Conflict of interests:

The authors declare not to have any interest conflicts.

Authors' contribution:

The authors have participated in the writing of the work and analysis of the documents



This work is licensed under a Creative Commons Attribution-NonCommercial 4.0 International license

Copyright (c) 2023 Edilio Aldana Pereira, Yordan Lores Pérez, Marisela Frías Tamayo

