

Chemical composition and *in vitro* digestibility of silages of taro (*Colocasia esculenta* (L.) Schott) tubers for feeding pigs

W. Caicedo¹, R. Rodríguez², P. Lezcano³, J. Ly³, S. Valle¹, L. Flores⁴ and F.N.A. Ferreira⁵

¹Secretaría Nacional de Educación Superior Ciencia Tecnología e Innovación, Quito, Ecuador

²Universidad de Granma, Centro de Estudios de Producción Animal, Bayamo, Cuba

³Instituto de Ciencia Animal, Apartado Postal 24, San José de las Lajas, Mayabeque, Cuba

⁴Escuela Superior Politécnica de Chimborazo, Riobamba, Ecuador

⁵Universidad Federal Minas Gerais, Belo Horizonte, Brasil

Email: orlando.caicedo@yahoo.es

To assess chemical composition and *in vitro* digestibility of silages of taro tubers (*Colocasia esculenta* (L.) Schott), for feeding pigs, four silages with waste tubers were carried out. A completely randomized design with a (4x6) factorial arrangement to analyze the chemical composition was applied. Natural yogurt (-1NY), whey (-2W), whey with 5% of molasses (-3WM5) and whey with 10% of molasses (-4WM10) were evaluated as silage components. The silages time were 15, 30, 60, 90 and 180 days. The *in vitro* digestibility of DM and OM with 15d of silages elaboration was investigated by means of simple classification design. The interactions treatments per day were significant ($P < 0.05$). The silage four in the 180th day show higher DM content (33.61%). The highest CP(N x 6.25) concentration (8.83%) was evidenced in silage two, in the 90th day. The lowest CF was determined in silage two in the 0(2.85) and 60th (2.84) days. The GE was high in silage one in the 0 day (4.40 kcal g DM⁻¹). The highest *in vitro* digestibility coefficient of DM (74.65 %) and OM (76.76 %) had it silage two. The conservation of taro tubers, between 0 and 180 d, by means of the addition of variables levels of natural yogurt, whey and sugar cane molasses, create different products with good DM, CP, GE content and low CF concentration, capable to be used in feeding pigs.

Key words: *by-products, fermentation, digestibility, preservation*

The efficient production of agricultural resources in tropical and subtropical areas, and the necessity to find, in a sustainable way, alternative sources for animal feeding are conditions that facilitate the use of silages in animal feeding (Guzmán 2010). By means of a simple and appropriate procedure, silages allows to preserve citrus residues (Llano *et al.* 2008), pineapple (Herrera *et al.* 2009), mango (Rego *et al.* 2010), between other wastes, useful for animals consume.

In Ecuador, according to Caicedo *et al.* (2013) there are a wide variety of feasible resources to feeding pigs, between them it is the taro (*Colocasia esculenta* (L.) Schott). Tubers are recognized as a cheap carbohydrates source of low cost, regarding to cereals and other cultivation types. They also present, high starch digestibility, that can reach up to 98% (Ezedinma 1987). The use of *in vivo* conventional methods to measure food digestibility and ingredients that are used in diets formulation is too expensive and requires long periods to obtain evaluative results. Consequently, have been developed *in vitro* methods that have the advantage of being simple, fast and allows evaluating a great number of samples at the same time and at low cost (Boisen and Fernández 1995 and Ramos 1995).

The objective of this study was to value chemical composition and *in vitro* digestibility of silages of taro (*Colocasia esculenta* (L.) Schott) tubers for feeding pigs

Materials and Methods

This study was carried out in the facilities of the

taro producer corporation of Pastaza, belonging to “Teniente Hugo Ortiz” parish, in Pastaza province, Ecuador. The weather in this area is semi-warm or humid sub-tropical, with precipitations between 4.000 and 4500 annually. It is located to an altitude of 950 m o.s.l, with average relative humidity of 87% and average minimum and maximum temperature of 18 to 26 °C. Soils are classified as inceptisols, oxisols and entisols (INAMI 2013).

To formulate the silos, Caicedo *et al.* (2013b) recommendations were followed (table 1). The micro-silos were prepared with tubers waste that, for its physical appearance, they do not fulfill the requirements established by national and international markets to use them in human feeding. Tubers were washed and milled in fresh way, in mix mill with blades and a sieve of 2.5cm, with the purpose of obtaining uniform particles.

For silages formulation, raw matters were weighed in a CAMRY digital balance, and were placed in four clean plastic tanks, with 400kg each. Ingredients were added in the following order: silage 1) cut tubers, natural yogurt and drinkable water for human consumption (1_NY) 2) cut tubers and whey (2_W); 3) cut tubers, molasses B (83° Brix) 5% and whey (3_WB5) and cut tubers, molasses B (83° Brix) 10 % (4_WB10).

The ingredients were mixed in a homogeneous way, manually, with a wooden spatula, during 15 min, at room temperature of 24 °C. Then, the silages mixtures were introduced in polyethylene bags, at a rate of 5kg. The bags were compacted and closed hermetically with

Table 1. Formulation of taro tubers silages

Raw matter inclusion, %	Treatments			
	1_NY	2_W	3_WMB5	4_WMB10
Cut taro tubers	68	68	68	68
Drinkable water for human consumption	27	-	-	-
Molasses B (83°Brix)	-	-	5	10
Natural yogurt	5	-	-	-
Whey	-	32	27	22
Total	100	100	100	100

1_NY: Silage with natural yogurt

2_W: Silage with whey

3_WMB5: Silage with whey and 5% of molasses B (83°Brix)

4_WMB10: Silage with whey and 10% of molasses (83°Brix)

a vacuum pump to guarantee their preservation. The micro-silos were stored under roof and protected of the sun light. They were opened at 0, 15, 30, 60, 90 and 180 d after silage.

The material was analyzed for DM, CF, and ash, CP (N x 6.25) content, SE and FNE, according to AOAC (2005). The levels of NDF, ADF and lignin were measured according to van Soest *et al.* (1991). It was considered that the OM concentration was equal to the difference 100- ash percent. However, the hemicellulose was the result of the subtraction NDF-ADF, as well as the cellulose (ADF-lignin). Both expressed in percent (van Soest and Robertson 1975). The GE was determined in a Parr adiabatic calorimetric pump, model 1241. The *in vitro* digestibility of the DM (DIMS) AND OM (DIMO) was carried out by pepsin-pancreatin-viscozyme, according to Boisen and Fernández (1991).

A completely randomized design with a (4x6) factorial arrangement for chemical characterization was applied, corresponding to four silage types (table 1) and six conservations time (0, 15, 30, 60, 90 and 180 d). *In vitro* digestibility was asses by means of a completely randomized design and analysis of simple variance was applied, with silage of 15 elaboration days. Three replications per each treatment were carried out. All determinations were made in triplicate. The means were contrasted by the analysis of variance technique, according to the recommendations of Steel *et al.* (1997). Where significant differences ($P < 0.05$) were found, they were compared with Duncan 1995) test. The analyses were carried out with the application of the Infostat statistical program ((Di Rienzo *et al.* 2012).

Results

During the conservation process, the average temperature was of 22 oC. When opening the micro-silos, the presence of alcohol or any symptom of ensilaged materials decomposition was never found. All of them had a sweet smell.

The result of the interactions studied for

chemical indexes is shown in table 2 and 3. The interactions treatments per day were significant ($P < 0.05$) for all chemical indexes evaluated: DM, OM, CP, CF, ash, SE, FNE, GE, NDF, ADF, lignin, hemicelluloses and cellulose.

The DM content was significantly ($P < 0.05$) higher in silage four in the 80th day (33.61%). While the highest OM concentration ($P < 0.05$) was determined in silage four in the 0 day (95.33%). The CF value ($P < 0.05$) was higher in silage two, in the 90th day (8.83 %). The lower CF content was determined in silage two in the 0 (2.85 %) and 60th (2.84%) days, respectively.

Silage four in the 0 day (4.67 %) showed the lower ash concentration. The higher SE content ($P < 0.05$) showed silage one, in the 30th (4.95 %) and 180th (4.95 %) days, consequently. The higher FNE concentration was evident in silage two, in the 30th (84.09 %) and 180th (84.09 %) days. The GE content was shown high in all the studied variants. However, was significantly ($P < 0.05$) high in silage one, in the 0 day (4.40 kcal gDM⁻¹).

The NDF was high ($P < 0.05$) in silage one in the 30th (18.21 %) and 180th (18.19 %) days. The lower ADF concentration ($P < 0.05$) have it silage two, in the 0 (2.55 %) and 60th (2.54 %) days. The higher lignin content ($P < 0.05$) was determined in silage four, in the 30th (2.46 %) day. The hemicelluloses decreased ($P < 0.05$) in silage four, in the 15th (9.48%) and 60th (9.48 %) days, and the cellulose increased ($P < 0.05$) in silage one, in the 30th (6.20%) and 180 th (6.19 %) days. The results of the digestibility study, IDDM and IDOM, of the four taro silages are showed in table 4. In the evaluation of DM and OM *in vitro* digestibility was showed that there were significant differences ($P < 0.05$).

The higher IDDM ($P < 0.05$) coefficient had it silage two (74.65 %). In silage one, the IDDM was lower (63.83%). Relate to IDOM, silage two showed the higher digestibility (76.76 %) coefficient ($P < 0.05$). Similarly, silage one had the lower digestibility coefficient (65.08 %).

Table 2. Chemical composition of silages of taro tubers. Interaction treatments x days (%in dry basis)

Indicator, %	Treatments	Days of ensilages					SE±sig	
		Day 180	Día 30	Día 60	Día 90	Día 180		
DM	1_NY	27.51 ^t	27.80 ^q	28.20 ^p	28.52 ⁿ	29.09 ^l	29.10 ^l	0.0227 P<0.0001
	2_W	27.60 ^s	28.20 ^p	28.51 ⁿ	29.32 ^k	29.55 ^j	30.23 ^f	
	3_MB5	27.71 ^r	28.33 ^o	29.80 ⁱ	30.09 ^g	30.68 ^e	31.25 ^d	
	4_MB10	27.82 ^q	28.91 ^m	29.91 ^h	31.92 ^c	32.94 ^b	33.61 ^a	
OM	1_NY	95.24 ^c	95.16 ^d	95.08 ^e	95.25 ^c	95.17 ^d	95.09 ^e	0.0075 P<0.0001
	2_W	93.69 ^h	93.58 ⁱ	93.45 ^j	93.69 ^h	93.59 ⁱ	93.46 ^j	
	3_MB5	93.38 ^k	93.33 ^l	93.12 ^m	93.39 ^k	93.33 ^l	93.13 ^m	
	4_MB10	95.33 ^a	94.23 ^f	94.13 ^g	95.29 ^b	94.24 ^f	94.14 ^g	
CP	1_NY	8.33 ⁱ	8.58 ^d	8.27 ^k	8.32 ⁱ	8.58 ^d	8.27 ^k	0.0031 P<0.0001
	2_SW	8.28 ^j	8.83 ^a	8.69 ^b	8.27 ^k	8.83 ^a	8.68 ^c	
	3_MB5	8.52 ^f	8.35 ^h	8.15 ^l	8.51 ^f	8.34 ^h	8.14 ^l	
	4_MB10	8.46 ^g	8.54 ^e	8.13 ^m	8.47 ^g	8.53 ^e	8.13 ^m	
CF	1_NY	3.35 ^k	5.78 ^c	5.85 ^b	3.35 ^k	5.78 ^c	5.85 ^b	0.0029 P<0.0001
	2_W	2.85 ⁿ	3.87 ^j	4.87 ^h	2.84 ⁿ	3.87 ^j	4.87 ^h	
	3_MB5	2.88 ^m	4.49 ⁱ	5.20 ^g	2.88 ^m	4.49 ⁱ	5.21 ^f	
	4_MB10	3.03 ^l	5.23 ^e	6.23 ^a	3.03 ^l	5.24 ^d	6.23 ^a	
Ashes	1_NY	4.76 ^k	4.84 ^j	4.92 ⁱ	4.76 ^k	4.83 ^j	4.92 ⁱ	0.0075 P<0.0001
	2_W	6.31 ^f	6.42 ^e	6.55 ^d	6.32 ^f	6.42 ^e	6.55 ^d	
	3_MB5	6.62 ^c	6.67 ^b	6.88 ^a	6.62 ^c	6.67 ^b	6.88 ^a	
	4_MB10	4.67 ^m	5.77 ^h	5.87 ^g	4.71 ^l	5.76 ^h	5.87 ^g	
SE	1_YN	4.72 ^d	4.88 ^c	4.95 ^a	4.72 ^d	4.89 ^b	4.95 ^a	0.0031 P<0.0001
	2_SL	2.77 ^j	2.83 ⁱ	2.83 ⁱ	2.77 ^j	2.83 ⁱ	2.84 ^h	
	3_MB5	3.26 ^g	3.42 ^f	3.53 ^e	3.26 ^g	3.43 ^f	3.53 ^e	
	4_MB10	2.17 ^l	2.16 ^l	2.46 ^k	1.93 ^m	2.17 ^l	2.45 ^k	
FNE	1_YN	78.85 ^p	81.88 ^f	83.88 ^b	78.84 ^p	81.86 ^g	83.86 ^c	0.0032 P<0.0001
	2_SL	79.79 ^o	81.09 ⁱ	84.09 ^a	79.78 ^o	81.08 ^j	84.09 ^a	
	3_MB5	78.72 ^q	78.55 ^r	80.55 ⁿ	78.72 ^q	78.55 ^r	80.56 ^m	
	4_MB10	81.31 ^h	80.73 ^l	82.73 ^d	81.32 ^h	80.75 ^k	82.72 ^e	
GE, kcalgDM ⁻¹	1_YN	4.40 ^a	4.23 ^d	4.23 ^d	4.23 ^d	4.23 ^d	4.23 ^d	0.0029 P<0.0001
	2_SL	4.21 ^e	4.26 ^b	4.19 ^f	4.26 ^b	4.18 ^g	4.19 ^f	
	3_MB5	4.23 ^d	4.26 ^b	4.26 ^b	4.25 ^c	4.26 ^b	4.25 ^c	
	4_MB10	4.25 ^c	4.19 ^f	4.25 ^c	4.19 ^f	4.25 ^c	4.25 ^c	

abcdefghijklmnopqrstuvwxyz Different superscripts show significant differences (P<0.05)

Table 3. Fibrous fractionation in silages of Taro tubers. Interaction treatments x days (% in dry basis)

Indicator, %	Treatments	Days of ensilages					SE±sig	
		Day 180	Día 30	Día 60	Día 90	Día 180		
NDF	1_YN	14.72 ^k	18.12 ^b	18.21 ^a	14.73 ^k	18.12 ^b	18.19 ^a	0.0083 P<0.0001
	2_SL	13.57 ⁿ	17.64 ^d	17.74 ^c	13.61 ^m	17.63 ^d	17.73 ^c	
	3_MB5	17.34 ^h	17.47 ^e	17.59 ^e	17.33 ^h	17.47 ^e	17.50 ^f	
	4_MB10	13.96 ^l	14.97 ^j	16.95 ⁱ	13.96 ^l	14.97 ^j	16.93 ⁱ	
ADF	1_YN	4.83 ⁿ	7.46 ^c	7.56 ^a	4.84 ⁿ	7.45 ^d	7.55 ^b	0.003 P<0.0001
	2_SL	2.55 ^r	6.34 ⁱ	6.84 ^e	2.54 ^r	6.34 ⁱ	6.83 ^f	
	3_MB5	4.18 ^o	6.27 ^k	6.57 ^h	4.17 ^o	6.25 ^l	6.56 ⁱ	
	4_MB10	4.07 ^p	5.49 ^m	6.62 ^g	4.05 ^q	5.49 ^m	6.62 ^g	
Lignin	1_YN	1.10 ^p	1.32 ^j	1.36 ^h	1.11 ^o	1.33 ⁱ	1.35 ^h	0.0035 P<0.001
	2_SL	1.12 ⁿ	1.21 ^m	1.28 ^k	1.13 ⁿ	1.21 ^m	1.27 ^l	
	3_MB5	0.81 ^s	1.71 ^f	1.81 ^e	0.82 ^r	1.69 ^g	1.82 ^e	
	4_MB10	0.83 ^r	2.07 ^c	2.46 ^a	0.84 ^q	2.05 ^d	2.43 ^b	
Hemicellulose	1_YN	9.90 ^j	10.66 ^b	10.65 ^h	9.89 ^j	10.67 ^h	10.65 ^h	0.0078 P<0.0001
	2_SL	11.02 ^e	11.30 ^b	10.90 ^g	11.07 ^d	11.29 ^b	10.90 ^g	
	3_MB5	13.16 ^a	11.21 ^c	11.02 ^e	13.16 ^a	11.22 ^c	10.94 ^f	
	4_MB10	9.89 ^j	9.48 ^k	10.33 ⁱ	9.91 ^j	9.48 ^k	10.32 ⁱ	
Cellulose	1_YN	3.72 ^k	6.15 ^b	6.20 ^a	3.72 ^k	6.12 ^c	6.19 ^a	0.0045 P<0.0001
	2_SL	1.43 ^r	5.13 ^e	5.56 ^d	1.42 ^r	5.14 ^e	5.56 ^d	
	3_MB5	3.37 ⁿ	4.56 ^h	4.73 ^f	3.35 ^o	4.56 ^h	4.74 ^g	
	4_MB10	3.24 ^p	3.42 ^m	4.16 ^j	3.21 ^q	3.44 ^l	4.19 ^j	

abcdefghijklmnopqrstuvwxyz Different superscripts show significant differences(P < 0.05)

Table 4. *In vitro* digestibility of DM and OM in silages of taro tubers

Variables	NY ⁻¹	SL ⁻²	MB5 ⁻³	MB10 ⁻⁴	EE ± sig
IDDM, %	63.83 ^d	74.65 ^a	70.86 ^c	74.49 ^b	0.0029 P < 0.0001
IDOM, %	65.08 ^d	76.76 ^a	72.15 ^c	76.07 ^b	0.0028 P < 0.0001

abcd Different superscripts show significant differences(P < 0.05)

Discussion

Chemical composition. The concentration of DM in the silages was in the recommended level (25-35 %) to characterized a silage of appropriate quality (McCullough 1975). In this regard, McDonald *et al.* (1981) suggested that 30% of DM was a minimum level to reduce the undesirable growth of clostridia that worse the product. The DM content of silages, until the 180th day of evaluation, were between 29.10 and 33.61%. Consequently, they can be considered good quality products for their use in animal feeding.

Regarding the CP content, the data of this research were in a concentration from 8.13 to 8.83%. Similar CP values informed Marrero *et al.* (1984) in silages of taro tubers of six months (8.5 %). Ogunlakin *et al.* (2012) referred higher results in tubers in natural way (4.93 to 5.17 %). They also made it Fetuga and Oluyemi (1976) in studies with cooked tubers (3.1 %). Apparently, by means of silage process the CP content is increased regarding

to tubers in natural and cooked way. The silages CP in the four variants showed low content (2.84 to 6.23%). In researches carried out with ensiled tubers, Marrero *et al.* (1984) determined 13.4% of CP. This wide variation in CP content could be due to the used ingredients in silages formulation. Also, the nutritional composition of roots and tubers vary according to climatic conditions, cultured varieties, soil conditions and time in which the crop is made (FAO 1990 and Onwueme 1999).

The FNE (4.67 to 6.88 %) and ash (4.67 to 6.88 %) concentration in silages showed higher content, regarding to that referred by Olajide *et al.* (2011) in fermented and dryings tubers. These authors informed 73.50 % for FNE and 0.88 % for ashes. Nevertheless, in relation to the SE content (2.63 %) referred by these researches, silages showed a similar performance to this nutrient (1.93 to 4.95%). The variation in FNE values could be due to the amount of fermented carbohydrates that the silages had in this experiment.

In this study, the GE content in silages was high, with values between 4.18 and 4.40 kcal gDM⁻¹ Abdurashid and Agwunobi (2009), in tubers in flour, informed 3.21 kcal g DM1, lower figures to those verified in this research. Apparently, the caloric content of ensiled tubers was increased by the energy contribution of natural yogurt, whey and molasses B.

In general lines, silages showed variation in NDF content (13.57 to 18.21%), ADF (2.55 to 7.56%), lignin (0.81 to 2.46%), hemicellulose (9.48 to 13.16%) and cellulose (1.42 to 6.19%). These variations can be associated to the lack or presence of molasses in the fermented material (Guzmán *et al.* 2012).

In vitro digestibility. In researchers developed by Ly and Delgado (2005), related with *in vitro* digestibility of DM and OM, with pancreatine pepsin, in fresh and dry taro tubers, informed that DM digestibility increased when tubers were dry (66.90%). Not happened this way when they were fresh (31.50 %). Likewise, the *in vitro* digestibility of OM was higher in the dry ones (76 %), in relation to the fresh ones (38.30 %).

It has been carried out several *in vitro* digestibility researches of DM and OM in other types of tubers in natural way. Ly *et al.* (2010) informed in cassava tubers values of 66 % of DM and 68.7 % of OM. In studies made with sweet potato, Ly *et al.* (1999) informed contents of 54.5 % of DM and 62.5 % of OM. These results were lower to those obtained in the study of the four silages variants. Apparently, the ensiled process and the digestibility method applied increased the *in vitro* digestibility coefficients of DM (63.83 to 74.65 %) and OM (65.08 to 76.76 %) consequently.

From the point of view of the OM *in vitro* digestibility, the ensiled taro tubers seem to be slightly better than other tropical tubers; due to in the ensiled process by means of anaerobic fermentation, lactic acid is produced and, in presence of this one, can be recover some wastes components, like protein, minerals and lipids (López *et al.* 2006).

The conservation of taro tubers, between 0 and 180d, by means of the addition of variables levels of natural yogurt, whey and sugar cane molasses originates different products that have good content of DM, CP, GE and low concentration of CF, suitable for its use in feeding pigs.

Acknowledgments

Thanks to the National Secretary of Higher Education, Science, Technology and Innovation (SENESCYT) of Ecuador for financing this research.

References

Abdurashid, M. & Agwunobi, L.N. 2009. Taro cocoyam (*Colocasia esculenta*) meal as feed ingredient in poultry. Pakistan J. Nut. 8:668
 AOAC. 2005. Official Methods of Analysis. 18th Ed. Association of Official Analytical Chemists, Inc. Gaithersburg, MD. USA.

Boisen, S. & Fernández, J.M. 1991. *In vitro* digestibility of energy and amino acids in pig feeds. In: Digestive Physiology in Pigs (M.W.A. Verstegen, J. Huisman y L.A. den Hartzog, eds.). Wageningen. Puduc. 231-236 pp.
 Boisen, S. & Fernández, J.A. 1995. Prediction of the apparent ileal digestibility of protein and amino acids in feedstuffs and feed mixtures for pigs by an *in vitro* analysis. Anim. Feed Sci. Tech. 51:29
 Caicedo, W.O., Rodríguez, R., Lezcano, P., Vargas, J., Ly, J. & Valle, S. 2013a. Efecto de inocuidad del ensilado biológico de tubérculos de papa China (*Colocasia esculenta* (L.) Schott) para la alimentación de cerdos. Revista Amazónica Ciencia y Tecnología 2:162
 Caicedo, W.O., Rodríguez, R., Lezcano, P. & Ly, J. 2013b. Estudios de composición química de ensilados de papa china (*Colocasia esculenta* L. Schott) destinados a la alimentación porcina. In: XXIII Reunión de la Asociación Latinoamericana de Producción Animal. La Habana. CD-ROM. ISBN 978 950 7171 492
 Di Rienzo, J., Casanoves, F., Balzarini, M., Gonzalez, L., Tablada, M. & Robledo, C. 2012. Grupo InfoStat, FCA. Universidad Nacional de Córdoba. Argentina. Versión 1.0 para Windows.
 Duncan, D.B. 1995. Multiple range and multiple F tests. Biometrics 11:1
 Ezedinma, F.O. 1987. Response of Taro (*Colocasia esculenta*) to water Management, Plot Preparation and Population. 3rd Intl. Symp. Trop. Root. Crops, Ibadan-Nigeria
 FAO. 1990. Roots, tubers, plantain and bananas in human nutrition. Effect of processing on nutritive values. Food and Agriculture Organization of the United Nations, Rome
 Fetuga, B.L. & Oluyemi, J.A. 1976. The metabolizable energy of some tropical tuber meals for chicks. Poultry Sci. 55:868
 Guzmán, O. 2010. Estudio de conservación de desechos de mango (*Mangifera indica* L.) para la alimentación de ovinos en el Estado de Nayarit. Master Thesis. Universidad Autónoma de Nayarit. Tepic. México. 87 pp.
 Guzmán, O., Lemus, C., Martínez, S., Bonilla, J., Plasencia, A. & Ly, J. 2012. Chemical characteristics of silages of mango (*Mangifera indica* L.) by products for animal feeding. Cuban J. Agric. Sci. 46:369
 Herrera, M., Chin-Jones, R. & Bourrillon, A. 2009. Características fermentativas y nutricionales del ensilaje del rastrojo de piña (*Ananas escamosus*). Agronomía Costarricense 33:15
 INAMHI. 2013. Instituto Nacional Meteorológico Hídrico. Reportes Climáticos del Ecuador. Pp. 20-25
 López, J., Sánchez, D. & Rosas, J. 2006. Analysis of free amino acid in fermented shrimp waste by high-performance liquid chromatography. J. of Chromatography A 1105:106
 Ly, J., Carón, M. & Delgado, E. 1999. A note on *in vitro* digestibility of sweet potatoes tubers (*Ipomea batata* (Lam) L.) for pigs. Cuban J. Agric. Sci. 33:179
 Ly, J. & Delgado, E. 2005. A note on *in vitro* (pepsin/pancreatine) digestibility of taro (*Xanthosoma sagittifolia* spp) and cocoyam (*Colocasia esculenta* spp) for pigs. Rev. Comp. Prod. Porcina 12:90
 Ly, J., Almaguel, R., Delgado, E., Carón, M. & Cruz, E. 2010. Estudios de digestibilidad *in vitro* (pepsina/pancreatina) de raíces de yuca para alimentar cerdos. Rev. Comp. Prod. Porcina 17:300
 Llano, D., López, D. & Mora, F. 2008. Potencial del ensilaje de desechos de naranja (*Citrus sinensis*). Rev. Ciencias

Técnicas Agropecuarias 17:41

- Marrero, L., Vargas, S. & Contreras, F. 1984. Ensilado de malanga japonesa (*Colocasia esculenta*) en la alimentación de cerdos en ceba. *Cienc. Téc. Agric. Ganado Porcino* 7:85
- McCullough, M.E. 1975. Nuevas tendencias en ensilaje de forrajes. *Rev. Mundial de Zootecnia* 15:44
- McDonald, P., Henderson, A.R. & Heron, S.J.E. 1991. *The Biochemistry of Silage*. Segunda edición. Marlow, Chalcombe. 340 pp.
- Ogunlakin, G.O., Oke, M.O., Babarinde, G.O. & Olatunbosun, D.G. 2012. Effect of drying methods on proximate composition and physic-chemical properties of cocoyam flour. *American J. Food Technol.* 7:245
- Olajide, R., Akinsoyinu, A.O., Babayemi, O.J., Omojola, A.B., Abu, A.O. & Afolabi, K.D. 2011. Effect of processing on energy values, nutrient and anti-nutrient components of wild cocoyam (*Colocasia esculenta* (L.) Schott) corm. *Pakistan J. Nut.* 10:29
- Onwueme, I. 1999. Taro cultivation in Asia and the Pacific. Food and Agriculture Organization of the United Nations. Regional Office for Asia and the Pacific. Bangkok, Thailand. 48 pp.
- Cuban Journal of Agricultural Science, Volume 49, Number 1, 2015
- Ramos, M. 1995. Aplicación de técnicas enzimáticas de digestión *in vitro* a la valoración nutritiva de piensos de conejos. PhD Thesis. C. Facultad de Veterinaria Departamento de Nutrición y Bromatología. Universidad Complutense de Madrid. Madrid. España.
- Rego, M.M.T., Neiva, J.N.M., Rego, A.C., Cândido, M.J.D., Carneiro, M.S.S. & Lobo, R.N.B. 2010. Chemical and bromatological characteristics of elephant grass silages containing a mango by-product. *Rev. Brasileira de Zootecnia* 39:81
- Steel, R.G.D., Torrie, J.H. & Dickey, M. 1997. *Principles and Procedures of Statistics. A Biometrical Approach*. MacGraw-Hill Book Company In Company. Third Ed. New York, USA. 666 pp.
- Van Soest, P.J. & Robertson, J.B. 1975. *Analysis of forages and fibrous foods*. Cornell University Press, New York, USA. p. 165
- Van Soest, P. J., Robertson, J. B. & Lewis, B. A. 1991. Methods for dietary fiber, neutral detergent fiber and nonstarch polysaccharides in relation to animal nutrition. *J. Dairy Sci.* 5:3583

Received: