

PODIUM

Journal of Science and Technology in Physical Culture

SCIENTIFIC PUBLICATIONS DEPARTMENT

Volumen 16
Issue 3

2021

University of Pinar del Río "Hermanos Saíz Montes de Oca"

Director: Fernando Emilio Valladares Fuente

Email: fernando.valladares@upr.edu.cu

Translated from the original in spanish

Original article

Comparison about the Jump capacity in junior athletes

Comparación de la capacidad de salto en deportistas juveniles

Comparaçãõ da capacidade de salto em atletas juvenis

Luis Benavides Roca^{1*}  <https://orcid.org/0000-0002-0995-2564>

Cristian Orellana Sepúlveda¹  <https://orcid.org/0000-0001-7978-735X>

Gastón Díaz Coria²  <https://orcid.org/0000-0002-3260-3071>

Hernán Maureira Pareja²  <https://orcid.org/0000-0002-4664-0287>

Anderson Dos Santos Carvalho³  <https://orcid.org/0000-0001-9928-3204>

Matías Morales González⁴  <https://orcid.org/0000-0001-5540-9725>

¹Universidad Santo Tomás. Chile.

²Catholic University of Maule. Chile.

³Universidade Paulista. Brazil.

⁴Université de Strasbourg. France.

*Corresponding author: benavides.roca@gmail.com

Received: 12/17/2020.

Approved: 27/07/2021.

How to cite ítem: Benavides Roca, L., Orellana Sepúlveda, C., Díaz Coria, G., Maureira Pareja, H., Dos Santos Carvalho, A., & Morales González, M. (2021). Comparison about the Jump capacity in junior athletes/Comparación de la capacidad de salto en deportistas juveniles. *PODIUM - Journal of Science and Technology in Physical Culture*, 16(3), 799-808. <https://podium.upr.edu.cu/index.php/podium/article/view/1053>



ABSTRACT

Within the dynamics of sports training the jumping evaluation is a useful tool to determine and compare the athlete's performance. The objective of the study is to compare the jumping ability between two groups of juvenile athletes, entering university in different years. The study is of a comparative type, 120 subjects were selected by convenience, which was distributed in two groups, according to the year in which they enter the university (2015 and 2020). Jumping capacity was measured with the 90° squat, countermovement and free squat protocols. The results showed a trend ($p=0.07$) in the countermovement modality, where the 2020 year group possessed better performance. In conclusion, the countermovement jump is projected as the comparison variable for populations with similar characteristics.

Keywords: Athlete; Motor performance; Physical Exercise.

RESUMEN

Dentro de la dinámica del entrenamiento deportivo la evaluación de salto es una herramienta útil para determinar y comparar el rendimiento del atleta. El objetivo del estudio es comparar la capacidad de salto entre dos grupos de deportistas juveniles, que ingresan a la universidad en diferente año. El estudio es de tipo comparativo, se seleccionaron 120 sujetos por conveniencia, los cuales fueron distribuidos en dos grupos, de acuerdo con el año en el que ingresan a la universidad (2015 y 2020). Se midió capacidad de salto con los protocolos de sentadilla de 90°, contra movimiento y libre. Los resultados mostraron una tendencia ($p=0,07$) en la modalidad de contra movimiento, donde el grupo del año 2020 poseía un mejor rendimiento. En conclusión, el salto en contra movimiento se proyecta como la variable de comparación para poblaciones de similares características.

Palabras clave: Atleta; El rendimiento del Motor; Ejercicio Físico.

INTRODUCTION

In high competition sports, there are sequences of explosive movements of short duration, which address an important part of the dynamics of the game (Ahmaidi, Buchheit, & Spencer, 2013), such as, for example, jumps, changes of direction, speeds and decelerations (Thomas *et al.*, 2020), these components involve mainly eccentric actions, capable of being reproduced as force situations linked to the success of the discipline (Harper *et al.*, 2020).

Different sport activities use the maximum mechanical power of the lower limbs as a reference point, to compare the quality and quantity of force produced by athletes (Jiménez *et al.*, 2009). Specifically, jumping in sport is considered a high-speed dynamic action, associated with the elevation of the body around an athletic target (Newton *et al.*, 2006). In this context, vertical lifting is a multi-joint action that demands levels of power and strength, which can be assessed through the 90° squat jump (SJ) and the countermovement jump (CMJ), which, in turn, are related to explosive strength and explosive elastic strength, respectively (Markovic, & Mikulic, 2010). In the same way, the magnitudes of strength achieved during this action are related to anthropometric components and sports performance (Bustos-Viviescas, Acevedo-Mindiola, & Merchán Osorio, 2020), therefore, it is very common to use this evolution within the sports field (Castagna *et al.*, 2013).



Based on the above, it is relevant to understand that the evaluation of jumping considers parameters of flight time, execution and take-off speeds, which are propitious data for training planning and determining the athlete's level (Bosco *et al.*, 1983; Di Giminiani, 2006), however, in the present study, only the height reached by the athlete will be addressed, due to the practicality generated by this variable and the greater possibility of comparison with the literature. Besides this, it allows projecting efficiency in mechanical and physiological processes, such as the production of tension, the recruitment of motor units and the frequency of nerve impulse discharge. Also, it represents the functioning of the elastic component of the muscle, which interacts with the motor control of the lifting technique (Saavedra *et al.*, 2017; Claudino *et al.*, 2017).

For its part, sports training leads to detailed knowledge of athletes, where evaluation turns out to be an essential tool within the progress of a subject. This allows comparing performances and establishing preparation structures that enhance the development of physical abilities (De la Rosa, *et al.*, 2020). Based on this premise, the objective of this research is to compare the jumping ability between two groups of youth athletes, who enter the university in different years, in order to identify the entry behaviors and establish the characteristics of each group according to the instances that were evaluated.

For this research some instruments were used, whose names and authors will be identified below:

- DM Jump brand contact platform (Saavedra and Vergara 2013).
- Methodology for evaluations (Bosco *et al.*, 1994).

MATERIALS AND METHODS

The present study is a cross-sectional study with a comparative design. As a participant, the sample was constituted with 120 subjects, belonging to the catchment program of the Universidad Católica del Maule (UCM), Chile. They were divided into two groups, according to the year in which they were evaluated (group A=2015, group B=2020). The athletes were part of the disciplines of soccer, volleyball, basketball, handball, athletics and gymnastics (Table 1).

Table 1. - Characterization of the sample

Year of income	n	Man	Woman	Old	Weight	Size
Group A 2020	44	29	15	19,5±1,3	71,5±11,5	172±0,0
Group B 2015	76	58	18	19,7±1,7	72,6±12	170±0,1
Total	120	87	33	19,6±1,5	72,1±11,8	171±0,0

The sampling used in this article is non-probabilistic by convenience. We proceeded to review the sports record and then performed the jumping evaluations.



Within the UCM protocols, the program for recruiting athletes for admission to the institution is established. Ethical guidelines and regulations are established, which must be taken into account at the moment of proceeding with the evaluations and the handling of the data; each athlete, prior to the evaluation, must sign an informed consent form where he/she declares to have knowledge of the characteristics of the tests. The above is governed by the Helsinki treaty and the ethical regulations of said university.

The measurements were taken on the DM Jump contact platform, where the data were processed using the DM Jump V2.2 Beta software, relating the flight time with the distance reached in the jumps.

Procedure

According to the jump evaluations based on the methodology announced in the introduction of this work, the SJ, CMJ and free modalities are treated. The protocol establishes to execute the SJ with a 90° knee flexion, with the feet in parallel and the hands on the hips. For the CMJ, the subject begins in a standing position with hands on hips, and then performs a 90° knee flexion and rises. As for the free jump, there is the consideration of executing it as a comfortable way as possible and trying to resemble some technical gesture of their sport. The time between each evaluation was approximately 3 minutes.

Prior to the evaluation, the athletes performed a 10-minute warm-up, which consisted of joint movements, dynamic actions and an adaptation to the technique of each jump.

Both groups followed the established protocol and the evaluations were carried out by the same researcher.

Analysis

Statistical analysis was performed in the SPSS Statistics 22 program. The data were subjected to a Shapiro-Wilk normality test. Descriptive statistics of mean and standard deviation were calculated. The t-test was performed for independent samples between the groups of athletes (2015 - 2020) considering a significance value $p \leq 0.05$ and a trend $p \leq 0.07$.

RESULTS

Figure 1 shows the jumping performance of the two samples of youth athletes, evaluated in 2015 and 2020. It is observed that there is no significant difference in the SJ ($p=0.11$) and free ($p=0.28$) modalities. While in the CMJ test it is possible to observe a trend between the two groups ($p=0.07$).






Fig. 1. - Comparison of jumps among athletes of the Universidad Católica del Maule

The present study investigated the performance of subjects belonging to the UCM recruitment program, which differed according to the instance in which they were evaluated (group A = year 2020 and group B = 2015). Therefore, the purpose of this article, lies in comparing the performance in jumping tests between the two samples of youth athletes, and thus, to be able to generate conclusions about the behavior of each group around the year in which they were measured.

DISCUSSION

The main finding of this research refers to the tendency shown by the CMJ test, which presents a higher performance of group A compared to group B. This is relevant for the analysis of sports performance, since this jump has a good prediction with the muscular power of the lower extremities (Markovic, *et al.*, 2004) and its technique possesses coordinative characteristics, which are related ($r = 0.9$) with the mechanical speed of discipline-specific gents (Linthorne, 2020). Along with this, Rodríguez-Rosell *et al.*, (2020) determined that actions associated with CMJ have speed as an enhancer factor, which stimulates neuromuscular processes, to achieve efficiency in the movement (+9.2%). Similarly, it has been seen that plyometric stimuli generate an increase in CMJ height (+0.6 %), which is due to the characteristics of the stimulus, since it favors muscle stretching and shortening actions in a sequential manner (Franco-Márquez *et al.*, 2015).

When comparing the results of the present study with the literature, it is observed that the sample of this research, has an inferior performance in the CMJ test to athletes who practice judo (43.3 ± 7.0 cm), jiu-jitsu (45.3 ± 4.1 cm), soccer (37.7 ± 7.4 cm), basketball (first division 45.51 ± 5.55 and second division 45.29 ± 6.13) and mixed martial arts (41.6 ± 6.1 cm) (Lima-Kons, *et al.*, 2017; Pehar, *et al.*, 2017; Campos, Leporce & Souto-Maior, 2019).



Regarding the results of SJ and CMJ, it can be seen that when contrasted with a group that practices BMX (bicycle motocross), at recreational level (CMJ= 48.7 ± 9.6 cm and SJ= 32.4 ± 8.1) and elite level (SJ= 50.9 ± 9.7 cm and 48.7 ± 9.6 cm) (Robert *et al.*, 2020), it is possible to identify that there is a lower jumping capacity. Similarly, when comparing the data with amateur athletes (SJ = 27.41 ± 7.6 cm and CMJ= 29.34 ± 8.30 cm), it is evident that group A has a higher performance, while group B, presents a lower flight height (Texeira *et al.*, 2019). In the same context, Yoshimura and Umemura (2016), evaluated a group of swimmers (SJ = 39 ± 7 and CMJ= 42 ± 7) and volleyball players (SJ= 45.5 and CMJ= 50 ± 5), which have a better performance than the athletes studied in this research.

When analyzing experimental studies, which show the improvement of jumping ability performance in a population of athletes, it has been seen that strength and coordination training, manages to improve the values of SJ (pre= 30.33 cm; post= 33.31 cm) and CMJ (pre= 31.9 cm; post= 35.25) in volleyball players (Kukiæ *et al.*, 2020). In this sense, Ramírez-Campillo *et al.*, (2016), evidenced an increase in flight height (SJ= 5.1%. and CMJ=4.4 %) in female soccer players, through a plyometric intervention. Falces-Prieto *et al.*, (2020), identified that U-19 soccer players increase their performance in the CMJ (pre= 34.4 cm; post= 37.1) with strength training with body weight load. According to these results, it is identified that UCM athletes have a lower performance, both before and after the intervention.

Among the limitations of the study is the lack of quantification of maturational parameters, which would have provided a higher level of interaction with jump height, since there is evidence that anthropometric aspects influence explosive actions such as jumps (Malina *et al.*, 2004; Almaiman, 2018).

CONCLUSIONS

In conclusion, the data of the present investigation show that there are no differences in the jumping ability in the groups of young athletes who enter the university in different years. Nevertheless, the CMJ evaluation could be the projectable test to obtain concrete results that identify the performance of UCM youth athletes.

Also, it is considered relevant, the quantification of the effect of the passage of five years, between two different groups, with similar contexts.

The use of jumps is recommended to determine the performance of the subjects and thus be able to generate training protocols that enhance the performance of youth athletes.



REFERENCES

- Ahmaidi, S., Buchheit, M., & Spencer, M. (2013). Confiabilidad, Utilidad y Validez de un Test de Sprint y Saltos Repetidos-BioKinetics. *Revista de Entrenamiento Deportivo*, 27(1). <https://g-se.com/confiabilidad-utilidad-y-validez-de-un-test-de-sprint-y-saltos-repetidos-1568-sa-H57cfb2722def>
- Almaiman, A. A. (2018). Effect of testosterone boosters on body functions: Case report. *International Journal of Health Sciences*, 12(2), 86-90. <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5870326/>
- Bosco, C., Komi, P. V., Tihanyi, J., Fekete, G., & Apor, P. (1983). Mechanical power test and fiber composition of human leg extensor muscles. *European journal of applied physiology and occupational physiology*, 51(1), 129-135. <https://pubmed.ncbi.nlm.nih.gov/6684025/>
- Bosco C. 1994. Test de Bosco. La valoración de la fuerza por el test de Bosco. Barcelona: Paidotribo.
- Bustos-Viviescas, B. J., Acevedo-Mindiola, A. A., & Merchán Osorio, R. D. (2020). Asociación de la masa muscular de miembros inferiores con el rendimiento en el salto vertical. *Revista Cubana de Investigaciones Biomédicas*, 39(3), e645. <http://www.revibiomedica.sld.cu/index.php/ibi/article/view/645>
- Campos Jr, J. C., Leporace, G., & Souto, A. (2019). Countermovement Jump Test Performance in Different Sports Modalities. *Journal of Exercise Physiology Online*, 22(5), 172-183. <https://go.gale.com/ps/i.do?id=GALE%7CA623324966&sid=googleScholar&v=2.1&it=r&linkaccess=fulltext&issn=10979751&p=HRCA&sw=w&userGroupName=anon%7E98d948e7>
- Castagna, C., Ganzetti, M., Ditroilo, M., Giovannelli, M., Rocchetti, A., & Manzi, V. (2013). Concurrent validity of vertical jump performance assessment systems. *The Journal of Strength & Conditioning Research*, 27(3), 761-768. <https://pubmed.ncbi.nlm.nih.gov/22648140/>
- Claudino, J. G., Cronin, J., Mezêncio, B., McMaster, D. T., McGuigan, M., Tricoli, V., & Serrão, J. C. (2017). The countermovement jump to monitor neuromuscular status: A meta-analysis. *Journal of science and medicine in sport*, 20(4), 397-402. <https://pubmed.ncbi.nlm.nih.gov/27663764/>
- De la Rosa, Y. A., Cajas, J. I. I., Pereira, L. G., & Rodriguez, Y. H. (2020). Herramienta de evaluación y su influencia en el proceso de formación del futbolista. *Revista Científica "Conecta Libertad" ISSN 2661-6904*, 4(2), 105-120. https://redib.org/Record/oai_articulo2819321-herramienta-de-evaluaci%C3%B3n-y-su-influencia-en-el-proceso-de-formaci%C3%B3n-del-futbolista
- Di Giminiani, R., & Scrimaglio, R. (2006). Center of gravity height calculation and average mechanical power during jump performance. *Italian Journal of Sport Sciences*, 13, 78-84. <https://es.scribd.com/document/44441552/Bosco-Jump-Calculations>



- Falces Prieto, M., González Fernández, F. T., Baena Morales, S., Benítez Jiménez, A., Martín Barrero, A., Conde Fernández, L., & Sáez de Villarreal, E. (2020). Efectos de un programa de entrenamiento de fuerza con autocargas sobre el rendimiento de salto con contramovimiento y la composición corporal en jugadores de fútbol jóvenes. *Journal of Sport & Health Research*, 12(1) 112-125. <https://dialnet.unirioja.es/servlet/articulo?codigo=7523032>
- Franco Márquez, F., Rodríguez Rosell, D., González Suárez, J. M., Pareja Blanco, F., Mora Custodio, R., Yáñez García, J. M., & González Badillo, J. J. (2015). Effects of combined resistance training and plyometrics on physical performance in young soccer players. *Int Sport Med*, 36, 906-914. <https://pubmed.ncbi.nlm.nih.gov/26180903/>
- Harper, D. J., Cohen, D. D., Carling, C., & Kiely, J. (2020). Can Countermovement Jump Neuromuscular Performance Qualities Differentiate Maximal Horizontal Deceleration Ability in Team Sport Athletes?. *Sports*, 8(6), 76. <https://www.mdpi.com/2075-4663/8/6/76>
- Jiménez, R., Parra, G., Pérez, D., & Grande, I. (2009). Valoración De La Potencia De Salto En Jugadores Semiprofesionales de Fútbol y Comparación De Resultados Por Puestos. *Cronos*. 14;79-84. <https://abacus.universidadeuropea.es/handle/11268/3261>
- Kukiaë, F., Todoroviaë, N., Èvoroviaë, A., Johnson, Q., & Dawes, J. J. (2020). Association of improvements in squat jump with improvements in countermovement jump without and with arm swing. *Serb J SportsSci* 11(1): 29-35. <https://www.iat.uni-leipzig.de/datenbanken/iks/power/Record/4061830>
- Lima Kons, R., da Silva Athayde, M. S., da Silva Junior, J. N., da Silva Tavares, W., & Detanico, D. (2017). Vertical jump performance in judo and Brazilian jiu-jitsu athletes: an approach with different training levels. *Ido Movement for Culture. Journal of Martial Arts Anthropology*, 17(4), 25-31. DOI:10.14589/ido.17.4.5 https://www.researchgate.net/publication/319460371_Vertical_jump_performance_in_judo_and_Brazilian_jiu-jitsu_athletes_An_approach_with_different_training_levels
- Linthorne, N. P. (2020). The correlation between jump height and mechanical power in a countermovement jump is artificially inflated. *Sports Biomechanics*, 1-19. <https://pubmed.ncbi.nlm.nih.gov/32200754/>
- Malina, R. M., Eisenmann, J. C., Cumming, S. P., Ribeiro, B., & Aroso, J. (2004). Maturity-associated variation in the growth and functional capacities of youth football (soccer) players 13-15 years. *European Journal of Applied Physiology*, 91(5-6), 555- 562. <https://pubmed.ncbi.nlm.nih.gov/14648128/>
- Markovic G.D., Dizdar I., Jukic M., & Cardinale M. (2004). Reliability and factorial validity of squat and countermovement jump tests, *Journal of Strength and Conditioning Research*, 18(3), 551-555. <https://pubmed.ncbi.nlm.nih.gov/15320660/>
- Markovic, G., & Mikulic, P. (2010). Neuro-musculoskeletal and performance adaptations to lower-extremity plyometric training. *Sports medicine*, 40(10), 859-895. <https://pubmed.ncbi.nlm.nih.gov/20836583/>



- Newton, R. U., Rogers, R. A., Volek, J. S., Häkkinen, K., & Kraemer, W. J. (2006). Four weeks of optimal load ballistic resistance training at the end of season attenuates declining jump performance of women volleyball players. *The Journal of Strength & Conditioning Research*, 20(4), 955-961. <https://pubmed.ncbi.nlm.nih.gov/17194257/>
- Pehar, M., Sekulic, D., Sisic, N., Spasic, M., Uljevic, O., Krolo, A., Milanovic, Z & Sattler, T. (2017). Evaluation of different jumping tests in defining position-specific and performance-level differences in high level basketball players. *Biology of sport*, 34(3), 263-272. <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5676323/>
- Ramírez-Campillo, R., González-Jurado, J. A., Martínez, C., Nakamura, F. Y., Peñailillo, L., Meylan, C. M., & Izquierdo, M. (2016). Effects of plyometric training and creatine supplementation on maximal-intensity exercise and endurance in female soccer players. *Journal of science and medicine in sport*, 19(8), 682-687. <https://pubmed.ncbi.nlm.nih.gov/26778661/>
- Robert, P., Cirer-Sastre, R., López-Laval, I., Matas-García, S., Álvarez-Herms, J., Julià-Sánchez, S., & Corbi, F. (2020). Relación entre capacidad de salto y rendimiento en ciclismo de BMX. *Apunts. Educación física y deportes*, 2(140), 37-43. <https://www.redalyc.org/journal/5516/551662868006/html/>
- Rodríguez-Rosell, D., Yáñez-García, J. M., Mora-Custodio, R., Pareja-Blanco, F., Ravelo-García, A. G., Ribas-Serna, J., & González-Badillo, J. J. (2020). Velocity-based resistance training: Impact of velocity loss in the set on neuromuscular performance and hormonal response. *Applied Physiology, Nutrition, and Metabolism*, 45(8):817-828. <https://pubmed.ncbi.nlm.nih.gov/32017598/>
- Saavedra, H. B., Medina, V. O., Mancilla, R. A., Pavez-Adasme, G., Da Silva, S. F., Hernández-Mosqueira, C., & Montoya, R. A. (2017). Fuerza de Salto Vertical en jugadores de Voleibol Varones de distinto nivel Competitivo. *Revista Horizonte Ciencias de la Actividad Física*, 8(1), 1-9. <http://revistahorizonte.ulagos.cl/index.php/horizonte/article/view/1>
- Teixeira, R. V., de Queiros, V. S., Dantas, M. P., Assis, M. G., Dantas, P. M. S., & Araújo Tinôco Cabral, B. G. (2019). Inter-limb performance asymmetry in amateur athletes involved in high intensity functional training. *Isokinetics and Exercise Science*, (Preprint), 1-7.
- Thomas, C., Dos' Santos, T., Comfort, P., & Jones, P. A. (2020). Male and female soccer players exhibit different knee joint mechanics during pre-planned change of direction. *Sports Biomechanics*, 1-14. <https://pubmed.ncbi.nlm.nih.gov/33115317/>
- Yoshimura, M., & Umemura, Y. (2016). Comparison of Factors Related to Jump Performance in Volleyball Players and Swimmers. *Journal of Exercise Physiology Online*, 19(4). <https://www.thefreelibrary.com/Comparison+of+factors+related+to+jump+performance+in+volleyball...-a0461945331>



Conflict of interests:

The authors declare not to have any interest conflicts.

Authors' contribution:

The authors have participated in the writing of the work and analysis of the documents.



This work is licensed under a Creative Commons Attribution-NonCommercial 4.0 International license
Copyright (c) 2021 Luis Benavides Roca, Cristian Orellana Sepúlveda, Gastón Díaz Coria, Hernán Maureira Pareja, Anderson Dos Santos Carvalho, Matías Morales González

